

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/257813444>

EEG-neurofeedback for optimising performance. I: A review of cognitive and affective outcome in healthy participants

Article in *Neuroscience & Biobehavioral Reviews* · October 2013

DOI: 10.1016/j.neubiorev.2013.09.015 · Source: PubMed

CITATIONS

195

READS

1,818

1 author:



John Gruzelier

Goldsmiths, University of London

402 PUBLICATIONS 13,256 CITATIONS

[SEE PROFILE](#)

Some of the authors of this publication are also working on these related projects:



Dissemination and cross fertilisation [View project](#)



Johrei and self hypnosis project, Imperial College London [View project](#)

Manuscript Number: NEUBIOREV-D-13-00251R1

Title: EEG-neurofeedback for optimising performance I: A review of cognitive and affective outcome in healthy participants.

Article Type: Special Issue: SAN

Keywords: neurofeedback; EEG; cognition; affect; validity; optimal performance; elderly

Corresponding Author: Prof. John Gruzelier,

Corresponding Author's Institution:

First Author: John Gruzelier

Order of Authors: John Gruzelier

Abstract: A re-emergence of research on EEG-neurofeedback followed controlled evidence of clinical benefits and validation of cognitive/affective gains in healthy participants including correlations in support of feedback learning mediating outcome. Controlled studies with healthy and elderly participants, which have increased exponentially, are reviewed including protocols from the clinic: sensory-motor rhythm, beta1 and alpha/theta ratio, down-training theta maxima, and from neuroscience: upper-alpha, theta, gamma, alpha desynchronisation. Outcome gains include sustained attention, orienting and executive attention, the P300b, memory, spatial rotation, RT, complex psychomotor skills, implicit procedural memory, recognition memory, perceptual binding, intelligence, mood and well-being. Twenty-three of the controlled studies report neurofeedback learning indices along with beneficial outcomes, of which eight report correlations in support of a meditation link, results which will be supplemented by further creativity and the performing arts evidence in Part II. Validity evidence from optimal performance studies represents an advance for the neurofeedback field demonstrating that cross fertilisation between clinical and optimal performance domains will be fruitful. Theoretical and methodological issues are outlined further in Part III.



Professor Verity Brown

27th September, 2013

Editor

Neuroscience & Biobehavioural Reviews.

Dear Verity

Further to your letter of today the manuscript entitled:

EEG-neurofeedback for optimising performance I: A review of cognitive and affective outcome in healthy participants.

Has been revised in the light of the reviewer's comments regarding minor corrections as follows: Points 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, and 17 have been corrected or clarified in line with suggestions.

Regarding Point 18 the section on alpha desynchronisation under the gamma beta section 5.1.2. has been deleted; it was already entered in the alpha section 5.1.4 and the duplication had been overlooked.

The reviewer is thanked for the helpful observations.

With best wishes,

John

Professor John Gruzelier

Professor Emeritus,

Department of Psychology

Goldsmiths

University of London



Professor Verity Brown

20th August, 2013

Editor

Neuroscience & Biobehavioural Reviews.

Dear Verity

The submission is the first of a three-part review on EEG-neurofeedback as a contribution to the SAN special issue:

EEG-neurofeedback for optimising performance I: A review of cognitive and affective outcome in healthy participants.

Parts II and III will be submitted within the next three weeks.

I suggest that Fabio Babiloni is the Guest Co-Editor that manages the reviews.

With best wishes,

John

Professor John Gruzelier

Professor Emeritus,

Department of Psychology

Goldsmiths

University of London



Professor Verity Brown

27th September, 2013

Editor

Neuroscience & Biobehavioural Reviews.

Dear Verity

Further to your letter of today the manuscript entitled:

EEG-neurofeedback for optimising performance I: A review of cognitive and affective outcome in healthy participants.

Has been revised in the light of the reviewer's comments regarding minor corrections as follows: Points 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, and 17 have been corrected or clarified in line with suggestions.

Regarding Point 18 the section on alpha desynchronisation under the gamma beta section 5.1.2. has been deleted; it was already entered in the alpha section 5.1.4 and the duplication had been overlooked.

The reviewer is thanked for the helpful observations.

With best wishes,

John

Professor John Gruzelier

Professor Emeritus,

Department of Psychology

Goldsmiths

University of London

- The first review of validity in EEG-neurofeedback optimal performance field.
- Over thirty controlled studies provide evidence of post-training benefits.
- Twenty-three in addition provide evidence of neurofeedback learning.
- Eight studies provide nine instances in support of feedback mediation in gains.

EEG-neurofeedback for optimising performance I: A review of cognitive and affective
outcome in healthy participants.

John H. Gruzelier, Goldsmiths, University of London

Correspondence address:

Professor John Gruzelier, Professor Emeritus, Department of Psychology, Goldsmiths
University of London, Lewisham Way, London SE14 6NW, UK.

Phone: 00 44 20 8 2992203

Email: j.gruzelier@gold.ac.uk

Keywords: neurofeedback; EEG; cognition; affect; validity; optimal performance; elderly

Abstract

A re-emergence of research on EEG-neurofeedback followed controlled evidence of clinical benefits and validation of cognitive/affective gains in healthy participants including correlations in support of feedback learning mediating outcome. Controlled studies with healthy and elderly participants, which have increased exponentially, are reviewed including protocols from the clinic: sensory-motor rhythm, beta1 and alpha/theta ratio, down-training theta maxima, and from neuroscience: upper-alpha, theta, gamma, alpha desynchronisation. Outcome gains include sustained attention, orienting and executive attention, the P300b, memory, spatial rotation, RT, complex psychomotor skills, implicit procedural memory, recognition memory, perceptual binding, intelligence, mood and well-being. Twenty-three of the controlled studies report neurofeedback learning indices along with beneficial outcomes, of which eight report correlations in support of a meditation link, results which will be supplemented by further creativity and the performing arts evidence in Part II. Validity evidence from optimal performance studies represents an advance for the neurofeedback field demonstrating that cross fertilisation between clinical and optimal performance domains will be fruitful. Theoretical and methodological issues are outlined further in Part III.

1.Introduction.

1.1. Historical Background.

Since the 1960's (Kamiya, 1968, 1969) the ability has been demonstrated for learned control of brain electrical activity through instantaneous feedback in the form of spectral power, event-related potentials or slow cortical potentials, putatively through operant conditioning or volitional control. This has now been extended to fMRI (e.g., Yoo et al, 2004; Weiskorf et al, 2004; deCharms, 2008; Rota et al, 2009; Hamilton et al, 2011; Zotev et al, 2011; Berman et al, 2012; Ruiz et al, 2013), transcranial doppler sonography (Duschek et al, 2012) and near infrared spectrometry (NIRS; Mihara et al, 2012; Kober et al, 2013). However, in contrast to autonomic and somatic peripheral biofeedback which became an established domain of Health Psychology (Feuerstein, 1988; Brannon and Feist, 2009), EEG-biofeedback was sidelined by science in the 1980s largely due to flawed studies and clinical overstatement, combined with the equivocal outcome of theoretically simplistic applications of alpha training for relaxation, when at the time psychophysiological understanding of alpha was at an early stage and was under the sway of unitary arousal models (Lindsley, 1952; Duffy, 1957). From the 1980s university research on EEG-biofeedback, later branded Neurofeedback, was confined to a few centres, notably under the direction of Sterman (e.g., Sterman and Friar, 1972; Sterman et al, 1974, 78; Sterman, 1996, 2000) and Lubar (e.g., Lubar and Shouse, 1976; Lubar and Lubar, 1984; Lubar et al, 1995a,b) in the USA and Birbaumer in Germany (Elbert et al, 1984; Rockstroh et al, 1989). Meanwhile a largely dedicated practitioner following arose outside of universities, especially in North America, encouraged by instrumentation companies. While helping keep the field alive and inventing diverse treatment protocols, this following mostly proceeded without scientific validation, sometimes accompanied by pseudo theorising and advertising speak.

Over the last decade or so university research, largely in Europe, has revisited neurofeedback, a revival of interest which the Society of Applied Neuroscience (SAN) has encouraged, and, as this review shows scientific evaluation and an evidence-base for applications is growing exponentially. This has come at a time when enhancing potential by ‘brain training’ is fashionable in contemporary culture (Owen et al, 2010; Rabipour and Raz, 2012), making the cause of scientific validation ever more essential and urgent, for aside from false promises, premature popularisation has led science to discard a field more than once.

1.2. Purview.

While ‘neurofeedback’ has now been applied to a range of brain imaging modalities, here the focus will be on the accumulating evidence of validation particularly this millennium in favour of neurofeedback applications with the EEG in healthy participants, commonly called the ‘optimal’ or ‘peak performance’ field. Validation has taken the form of controlled studies showing differential group outcomes advantaging a neurofeedback protocol, and more importantly has included correlation between the feedback learning and the outcome allowing the mediation of neurofeedback learning to be inferred. The review will be initially structured around neurofeedback protocols in conjunction with psychological processes, followed by applications in sport and with the elderly. A companion review, Part II, will cover applications in the performing arts and creativity (Gruzelier, 2013a) while Part III will be devoted to methodological and theoretical issues in order to help advance this re-emergent field (Gruzelier, 2013b). The review will not concern clinical applications which have been the main concern of the field to date. To give some examples there are reviews on Attention Deficit Hyperactivity Disorder (ADHD; Arns et al, 2009; Lofthouse et al, 2012), epilepsy (Tan, 2009), autism spectrum disorder (Coben et al, 2009); controlled studies on addiction (Scott et al, 2005), tinnitus (Hartmann et al, 2013) and insomnia (Cortooos et al, 2010); case

1 studies on stroke (Bearden et al, 2003), anxiety and depression (Hammond, 2005) and
2 cognitive disorganisation with anhedonia in conjunction with drug misuse (Unterrainer et al,
3 2013). Similarly there are reviews on the burgeoning approach branded Brain Computer
4 Interface (BCI), one typically focussing on the neuro-rehabilitation of patients with
5 tetraplegia from brain or spinal cord injuries enabling patients through EEG feedback to
6 communicate and/or physically interact with their environment, and up until now largely
7 consisting of engineering innovations and a small clinical evidence base (Birbaumer et al,
8 2008; Mak and Wolpaw, 2009; Silvoni et al, 2011).

1.3. Protocols.

As will be seen a diversity of neurofeedback training protocols has been applied for
optimising performance. The most popular one has involved training-up the amplitude of the
Sensory Motor Rhythm (SMR) 12 – 15 Hz band while inhibiting outer-lying bands in the
EEG spectrum. This had followed evidence that within the 12-15 Hz frequency range
recordings from sensory-motor and pre-motor cortices showed a distinctive oscillation which
was maximal during periods of quiet wakefulness, with reduced muscle tone, and was absent
during goal directed activity and desynchronising with motor intention (Serman and
Wyrwicka, 1967; Wyrwicka and Serman, 1968; Serman and Friar, 1972; Serman et al,
1974). Notably application of the SMR neurofeedback protocol has reduced motor seizure
rates in epileptic patients while normalising their sleep patterns and EEG (Serman, 2000;
Serman and Friar, 1972; Serman et al, 1974, 1978). By extrapolating to ADHD the potential
of reducing the excitability of the sensorimotor system with concomitant suppression of theta,
and following SMR training with training up adjacent low beta activity (16 – 22 Hz; beta1)
an index of EEG desynchronisation, improvements in attention and hyperactivity were first
demonstrated in case studies and early controlled trials (e.g., Lubar and Shouse, 1976; Lubar

et al, 1995a; Rossiter and La Vaque, 1995; Linden et al, 1996) and now have a substantive evidence base (Monastra et al, 2005; Arns et al, 2009; Lofthouse et al, 2010).

Another pioneering protocol involved raising the theta-alpha ratio with auditory feedback and eyes closed, termed alpha/theta (A/T) training. Following on from the earliest attempts to up-train alpha for relaxation and reduce anxiety (Budzynskie and Stoyva, 1972; Hardt and Kamiya, 1978), the focus on theta first grew out of diverse cultural evidence that the deeply relaxing, hypnagogic reverie or twilight theta state was conducive to creative insights (see Gruzelier, 2009). From clinical studies the goal evolved of elevating theta over alpha to achieve crossover. Early on this had demonstrable benefits for addiction and PTSD which included enhanced well-being when introduced as a central part of a therapeutic package (e.g. Peniston and Kulkowsky, 1989; 1991; Saxby and Peniston, 1995).

Subsequently the SMR and A/T protocols have been contrasted for their optimal performance effects in healthy participants, while new neurofeedback protocols have been developed from contemporary neuroscience including training upper-alpha, peak alpha frequency, gamma, and various theta protocols, the subject of this review. Historically self-regulation of slow cortical potentials (SCPs) had received extensive validation (Elbert et al, 1984; Rockstroh, 1989), but has not attracted interest in the optimal performance field despite promise in trials with ADHD where it has been compared favourably with EEG-spectrum training (Gevensleben et al 2009).

1.4. Issues Considered.

In what must be a necessarily concise selection of study features to be considered the following issues have been documented.

What evidence is there that feedback learning occurred? Have learning indices been reported or is learning inferred from demonstration that a group receiving neurofeedback outperformed a comparison group? What would evidence of learning consist of: learning curves

1 within sessions, across sessions, the correlation of learning indices with outcome
2 assessments? If the most basic theoretical premise about neurofeedback holds, namely that
3 the brain's rhythms are changed by training, is learning reflected in subsequent training
4 sessions including baselines?
5
6
7

8
9
10 There are issues of specificity. i) Is there band specificity or independence such that
11 only the trained bands are influenced or is there leakage or even reciprocity within the EEG
12 spectrum? ii) Is there specificity and independence regarding cognitive/affective outcome
13 such that performance enhancement is specific to some processes leaving other processes
14 unchanged? iii) Is there topographical specificity such that the EEG outcome is specific to the
15 training site, or is it distributed locally beyond the training site, or is it only distal from the
16 training site?
17
18
19
20
21
22
23
24
25

26
27 Are there sufficient sessions to give learning a chance given that a learning process is
28 being built? Can learning even be obtained from one-session and if so what is the validation?
29 What is the interval between sessions given there are considerations for learning of spaced
30 versus distributed practice. Are some people unable to learn to self-regulate their brain
31 rhythms – sometimes termed non-responders? The outsider may be surprised at how few
32 subjects are typically included in most intervention studies. Consider though that with say
33 fourteen subject sessions to include a ten session course of training plus two pre and post-
34 training outcome assessments, then for a study involving three groups of ten this would add
35 up to 420 experimenter sessions to acquire data, and not allowing for dropout replacements
36 which are inevitable when enterprising, mostly student subjects are required to fit fourteen
37 sessions into their busy lives; let alone the logistics of engaging as trainees professionals such
38 as eye surgeons (Ros et al, 2009).
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55

56 All these issues will be reconsidered in detail in Part III focussing on methodology and
57 theory (Gruzelier, 2013b). In Part I the over-riding objective will be to weigh up the now
58
59
60
61
62
63
64
65

1 rapidly accumulating evidence base in the search of validation of neurofeedback, a concern of
2 the author's from an early review (Gruzelier et al, 2006). As will be seen the majority of
3
4 studies report evidence of neurofeedback learning and in this respect the optimal performance
5
6 field is far ahead of other neurofeedback domains in reporting evidence of learning.
7
8
9

10 11 12 2. EEG Spectral Neurofeedback.

13 14 2.1 SMR and Diverse Protocols.

15 16 2.1.1. Sustained Attention, Selective Attention and Memory: SMR and Beta1 Training.

17
18 SMR and beta1 neurofeedback protocols widely used with ADHD were found to have a
19
20 positive impact on the sustained attention of healthy subjects in two studies (Egner and
21
22 Gruzelier, 2001; 2004a) which were an adjunct to music performance enhancement (Egner
23
24 and Gruzelier, 2003; see for review Gruzelier, 2012, Gruzelier, 2013a). In the first music
25
26 study as a secondary aim SMR (12-15 Hz, C4) and beta1 (15-18 Hz, C3) protocols with theta
27
28 (4-7 Hz) and high beta (22-30 Hz; beta2) inhibits were compared in twenty-two music
29
30 students. Protocols were administered each for 15-mins, twice a week for five weeks. A
31
32 timed continuous performance, go / no-go, attention task (Greenberg and Kindschi, 1999),
33
34 used as a gold standard in the ADHD field (Monastera et al, 2005; Arns et al, 2009), assessed
35
36 attentiveness through omission errors and impulsiveness through commission errors while a
37
38 perceptual sensitivity or d-prime index took into account both error types. Additionally the
39
40 auditory event-related potential P300b was recorded with a 28-electrode montage, along with
41
42 the resting EEG (Egner et al, 2004).
43
44
45
46
47
48
49
50

51 The behavioural task disclosed a significant reduction in commission errors
52
53 contributing to an increase in d-prime, which were positively associated in regression
54
55 analysis with learned increases within session in SMR and beta1 amplitudes, especially SMR
56
57 amplitudes, learning indices that were positively correlated ($r = 0.65$, $p < 0.01$). When the
58
59
60
61
62
63
64
65

1 effect of beta1 on SMR was examined with partial correlation, the positive relation between
2 SMR and commission error reduction was increased ($r = -0.79$, $p < 0.001$), and see Figure 1,
3 while when controlling for the effect of SMR on beta1 the opposite relation with the beta1
4 increment was disclosed ($r = 0.71$, $p < 0.001$) in keeping with increased arousal. Therefore
5 evidence was provided of protocol specific effects on the behavioural measures of attention,
6 in other words SMR and beta1 training had different consequences and were not
7 interchangeable and importantly supported a mediational link between the SMR training and
8 increased attentiveness.
9

10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
Figure 1 about here.

The psychophysiological assessment disclosed that event-related P300b amplitudes following training were larger across frontal (F3,FZ, F4), central (C3, CZ, C4) and parietal (P3, PZ, P4) scalp, with the SMR effect stronger with frontal and C3 derivations and the beta1 effect stronger with central derivations. Otherwise the relations between the protocols and the P300b did not differ for both protocols disclosed positive correlations with learning indices (SMR, $r = 0.49$, $p < 0.06$; beta1, $r = 0.55$, $p < 0.05$, and see Figure 2) in support of common influences on perceptual and cognitive integration processes reflected in the P300b, possibly through top-down resource allocation. In contrast the opposite relations of the protocols with behavioural attention measures conceivably may reflect converse influences on the subsequent motor response as required by the timed go /no-go sustained attention task (Egner and Gruzelier, 2001).

Figure 2 about here.

A second study (Egner and Gruzelier, 2004a) involved an independent group design with another group of twentyfive conservatoire students randomised to ten weekly sessions of SMR and beta1 (Cz) training or to a control group trained in the Alexander technique popular in performing arts conservatoires and involving postural and stress reduction. The sustained

1 attention task and auditory P300b paradigm were repeated and supplemented by a more
2 complex auditory divided attention task. Both behavioural attention tasks showed an increase
3 in d-prime following SMR training, here attributable to improvements in commission errors
4 as well as omission errors in the more complex task. Response time variability in the divided
5 attention task was also improved in the SMR group, while a single advantage in the beta1
6 group was a reduction in response time. These results provided further evidence of protocol
7 specificity, as was the case here with the P300b amplitudes which were enlarged following
8 beta1 training only especially at central and parietal placements. A topographical resting eyes
9 closed EEG outcome assessment with both studies was also undertaken which mainly
10 disclosed a frontal long-term locus (Egner et al, 2004) and is discussed in Part III.
11
12
13
14
15
16
17
18
19
20
21
22
23

24 The data from the two studies were interpreted as indicating a general attention-
25 enhancing effect of SMR training beyond impulse control, and an arousal-enhancing effect of
26 beta1 training. Theoretically the SMR result was consistent with improved response
27 inhibition following inhibition of sensory-motor cortex (Serman, 1996), while beta1
28 enhanced the signal-to-noise ratio in stimulus processing. The latter when combined with an
29 impulsive response tendency was in keeping with heightened activity in a noradrenergic
30 vigilance network, one associated with fast but commission error-prone response tendencies
31 in models of attention (Posner and Raichle, 1996). The differential group effects and the
32 correlations with feedback learning discounted nonspecific influences such as generic
33 learning effects, practice, motivation and expectancy and so validated the impact of
34 neurofeedback learning.
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50

51 In a third study (Vernon et al, 2003) with a primary interest in training up theta in order
52 to improve working memory (see 2.4.2) medical students were randomised to one of three
53 groups: SMR training while inhibiting theta (4-7 Hz) and beta (18–22 Hz), eyes open theta
54 training while inhibiting delta (0-4 Hz) and alpha (8 – 12 Hz), both with a Cz placement, and
55
56
57
58
59
60
61
62
63
64
65

1 a non-training control group. The effect of eight twice-weekly sessions were examined on a
2 sustained attention task with memory load varied by a two- or three-digit sequence target, and
3
4 on a cued-recall semantic working memory task with words presented in semantic clusters or
5
6 randomly (Haarman et al, 2003). While there was clear evidence of operant control over the
7
8 SMR from a within-session increase in SMR and reductions in theta and beta, this was not
9
10 achieved with eyes-open theta training and indeed alpha increased relative to theta (2.5.2).
11
12 Following SMR training the two-digit CPT disclosed highly significant reductions in errors
13
14 of omission and commission which were not observed in the other groups, whereas outcome
15
16 on the three-digit task with it greater demands on memory load was not improved by either
17
18 protocol. Invoking a three-network model of attention (Posner and Peterson, 1990) it was
19
20 inferred that SMR training may act on alerting and orienting networks favouring simple
21
22 target detection as in the 2-digit task and not the executive attention network invoked in the
23
24 3-digit task with the greater memory load; though with hindsight eight sessions may have
25
26 been insufficient for the more difficult task.
27
28
29
30
31
32
33

34 Notwithstanding, the more robust effects in favour of SMR training were found with
35
36 the semantic working memory task where there were gains of the order of 10% with both
37
38 clustered and unclustered recall. Working memory has been shown to involve the interaction
39
40 of attention and memory storage circuitry including respectively a frontal executive control
41
42 system and posterior association cortex (Sarnthein et al., 1998; von Stein et al., 1999; von
43
44 Stein and Sarnthein, 2000), while Haarmann and Cameron (2005) have reported an increase
45
46 in coherence in the 10–14 Hz band between frontal and posterior regions during semantic
47
48 working memory. This connectivity may be facilitated by SMR (and upper-alpha) training.
49
50
51
52
53

54 The sustained attention task (Greenberg and Kinschi, 1999) has also been examined in
55
56 11 year olds as an adjunct to a creative music performance study (Gruzelier et al, 2013;
57
58 Gruzelier, 2012, and see Part II, Gruzelier, 2013a). Thirty-three children were randomised to
59
60
61
62
63
64
65

10 weekly sessions of an SMR/theta/beta2 protocol, or to A/T training, or to a no-intervention control group. Cumulative learning was found within and across sessions for theta/alpha and SMR/beta2 ratios, while the SMR/theta ratio remained unchanged. Referring here only to the outcome with sustained attention, there was a highly significant improvement in d-prime for the neurofeedback groups, largely attributable to a reduction in commission errors. The gain was much greater following A/T training with only a weak improvement following SMR training, however, in contrast to the increase in the SMR/ beta2 ratio the SMR/theta ratio was unchanged indicating less than optimal learning. That these benefits, which aside from sustained attention included improved music performance and well-being (Gruzelier et al, 2013a), occurred with only 10 sessions of training has important pedagogic implications for the value of neurofeedback in school children, quite aside from the fact that according to the attention test manual 19/33 children were classifiably ADHD on the basis of their attention performance and may not have been expected to respond with only 10 sessions when in the clinic thirty would be a minimum.

In children of a similar age, and comparing girls with boys, Barnea et al (2005) examined a Hebrew word/nonword recognition decision task allowing an independent contribution to reading from each hemisphere. 20 Israeli 10-12 year olds received SMR/theta training for 20 half hour sessions at either C3 or C4, with groups counterbalanced for gender. Setting aside results that in the absence of a control group may have reflected possible practice or nonspecific confounds, the protocols improved language processing in the hemispheres differentially – C3 training improved word recognition and C4 nonword recognition, in line with left hemispheric specialisation for words and the right for nonwords. At the same time whereas accuracy of reading was improved by training at C3 in girls, it was improved by C4 training in boys from which the authors inferred the involvement of contralateral callosal circuitry. Bilateral circuitry was also implicated, for whereas latency of

1 response to nonwords was improved with C4 training, C3 training improved both word and
2 nonword recognition. In the authors' words: "The results suggest that the lateralized NF
3
4 protocol activates asymmetric hemispheric control circuits which modify distant hemispheric
5
6 networks and are organized differently in boys and girls." p 314.
7
8

9 2.1.2. Visuo-Motor Skills: SMR & Alpha/Theta. 10

11 SMR/theta and A/T neurofeedback protocols have also been applied to enhance the
12 visuo-motor skills of junior doctors training to become ophthalmic surgeons (Ros et al,
13
14 2009). Twenty doctors were randomised to either SMR ratio or theta/alpha ratio protocols, of
15
16 whom 8/20 before hand participated in a wait-list control group. Outcome was measured by a
17
18 simulated cataract surgery skills lab procedure assessed by timed measures, self-report
19
20 anxiety ratings and technique ratings from filmed performance by two consultant surgeons
21
22 (SMR group inter-rater reliability, $r = 0.85$). While improvements following A/T training
23
24 were not significant and there was no change in controls, following SMR training the time on
25
26 task was improved by 26% and this was greatest for the more complex suture task. The gain
27
28 following SMR training with the objective timed measures found a counterpart in the
29
30 subjective consultant ratings of improvement especially for the suture task, and furthermore
31
32 the timed improvement correlated with the surgeon ratings. The SMR group also showed a
33
34 reduction in anxiety.
35
36
37
38
39
40
41
42

43 Figure 3 about here.
44

45 Further light was shed through positive relations between within- and between-session
46
47 learning from both protocols and microsurgical performance. Interestingly within-session
48
49 SMR learning correlated with the pause time between tasks ($r = 0.72$, $p < 0.019$) when
50
51 allowing with partial correlation for the association between longer pause time and more
52
53 efficient performance overall ($r = 0.70$, $p < 0.035$). As the pause between tasks was longer
54
55 with SMR training, though not significantly so, this implied that SMR training in producing a
56
57
58
59
60
61
62
63
64
65

1 better performance achieved this through a more modulated performance, faster performance
2 overall with longer, perhaps preparatory pauses between tasks. Though A/T training did not
3 significantly improve performance overall, the better the within-session A/T learning the
4 greater was the improvement in surgical technique ($r = 0.64$, $p < 0.047$) and the more efficient
5 was the technical performance ($r = -0.52$, $p < 0.06$). When the A/T group was median split into
6 those with the higher and lower improvement in technique, it was the higher improvement
7 group who showed the superior theta/alpha ratio, an index of hypnogogia (Gruzelier, 2009).
8 Of possible relevance to these individual differences was the challenge of achieving the A/T
9 goal of borderline sleep within sessions given that the half hour sessions were sandwiched
10 into the busy schedules of the junior doctors. An increase in the theta/alpha ratio was also
11 associated with a reduction in performance anxiety. A complimentary result was found when
12 the same median split was applied to the SMR group; those in the lower improvement group
13 showed a 10% falloff in the SMR/theta ratio, shown to be attributable to an increase in theta
14 in later sessions, the opposite to the training goal of inhibiting theta. Poorer improvement in
15 surgical technique was also found to be accompanied by a greater lapse of time between
16 training sessions, an important methodological issue (Gruzelier, 2013b). In conclusion both
17 SMR and A/T could be seen to benefit psychomotor skills when effective learning was taken
18 into account.

2.1.3. Visuospatial Rotation and RT: SMR Ratio and Theta/Beta1 Reduction.

21 Doppelmayer and Weber (2011) delivered thirty training sessions (C3/4) to an SMR
22 (12 – 15 Hz) group ($N = 13$) or to a group ($N = 14$) receiving a theta (4.5 – 7.5 Hz)/beta1 (15
23 – 21 Hz) ratio reduction protocol which has had a successful outcome in ADHD research
24 (Leins et al, 2007; Gevensleben et al, 2009). There were theta, beta2 and eye-blink inhibits.
25 In addition a control group ($N = 14$) was instructed to either increase or decrease the
26 amplitude of different 1-Hz bands within the range of 6 to 35 Hz, variable over sessions.

1 Only the SMR group evinced learning, showing a highly significant progressive linear
2 amplitude increase accompanied by a significant increase in reward threshold; though
3
4 learning only differed from baseline after 26-30 days of training. Pre-and post-training
5
6 assessments included spatial rotation, simple and choice reaction time, letter cancellation and
7
8 a creativity test. Of these measures SMR training facilitated spatial rotation and simple and
9
10 choice reaction time, tasks which required visuospatial processing, access to semantic
11
12 memory, and the integration of relevant stimuli. SMR training benefits were attributed to
13
14 improved regulatory control of somatosensory and sensorimotor pathways, which in turn led
15
16 to more efficient attention resulting in a better integration of task-relevant stimuli, in line with
17
18 Egner and Gruzelier, (2001).
19
20
21
22
23

24 Figure 4 about here.
25
26

27 2.1.4. Sleep, Memory and SMR. 28 29 30

31 A series of studies has shown that SMR training without inhibits transfers to the
32
33 sleeping state. The potential for SMR training to enhance sleep arose from the earliest
34
35 demonstrations in cats (Serman et al, 1969) following evidence that a dominance of
36
37 oscillations overlapping the SMR 12 – 15 Hz band occurred in stage II NREM sleep, leading
38
39 to the descriptor of sleep spindles (rolandic mu or wicket rhythms; Niedermeyer, 2005).
40
41 Cognitive associations with sleep spindles in recent reports have included over-night memory
42
43 consolidation of motor procedural memory and declarative memory, learning aptitude,
44
45 implicit learning and intelligence (e.g., Fogel et al, 2007; Shabus et al, 2004). Hauri et al
46
47 (1981,1982) went on apply SMR training in the treatment of insomnia in controlled studies.
48
49
50
51
52
53

54 Berner et al (2006) in a pilot study with healthy subjects explored the effect of Cz
55
56 neurofeedback in the range 11.6–16 Hz on sleep spindle activity (C3,C4) and the sleep EEG
57
58 (Fz, C3, Cz, C4, Pz, O1, O2) while declarative memory was assessed after neurofeedback and
59
60
61
62
63
64
65

again after a night in the sleep laboratory. Eleven participants previously shown to be capable of raising their alpha power through feedback were assigned either to neurofeedback or to the same protocol but with sham feedback. The protocols were given in a single session in four ten-minute training blocks each separated by 2 minute breaks. While the experimental group was successful in raising power in the training band, the single neurofeedback session did not have an impact on either spindle activity in sleep or declarative memory. However, though compromised by small group sizes a tendency was reported for power in the training band to be increased at Fz and Cz relative to the pre-training baseline in the first half of the night in stage 2-4 NREM sleep, a period during which the spindle activity correlated positively with memory. When all eleven subjects were categorised according to their overall memory those with the higher memory score showed more spindle activity during the early period of sleep in support of an aptitude to learn new verbal items (Gruber et al, 2006). The resting eyes-open EEG was examined for carryover effects immediately after neurofeedback, and while there was no increase in sigma the authors reported ‘a noticeable’ but statistically nonsignificant increase in slower 4-10 Hz bands, which would be in keeping with relaxation.

Hoedlmoser et al (2008) went on to assign twenty-seven participants to 10 sessions of either SMR (12 – 15 Hz) C3 training without inhibits or to a control group receiving randomised frequency training of 3 Hz bins between 3 and 20 Hz, excluding the SMR band. Pre- and post-training assessment consisted of a declarative word-pair memory task before and after a 90 minute afternoon nap in the sleep laboratory, as well as EEG sleep characteristics from a 15 electrode recording. In addition to learned changes during feedback sessions, the impact of neurofeedback was also estimated from successive pretraining baselines with the assumption of cumulative learning carrying over day by day. There was evidence of successful learning through an increase in SMR amplitude from early to late sessions, and while a cumulative increase across SMR baselines was not observed there was a

1 transfer to the tonic EEG in the post training nap with an increased spindle number and
2 shorter sleep latency indicative of improved sleep quality. Furthermore following SMR
3 training immediate retrieval in the declarative memory task was facilitated both before and
4 after the nap, whereas neither group showed improvements in memory consolidation as a
5 result of the nap per se. This was interpreted as neurofeedback improving memory through an
6 indirect nonspecific influence facilitating attention or relaxation during learning/retrieval of
7 the task.
8
9

10
11
12
13
14
15
16
17 SMR training was later extended by the same group (Schabus et al, 2013) to twenty-
18 four patients with primary insomnia who in a counterbalanced design each received 10
19 sessions of SMR C3 training together with 5 sessions of sham training involving random 3
20 Hz bins between 7 – 20 Hz excluding SMR, which were scheduled over 3 – 6 weeks. SMR
21 was monitored bilaterally along with beta (16 – 25 Hz). At C3 there was a linear increase
22 during SMR training in both SMR and broad-band beta amplitude, with a similar but
23 nonsignificant effect contralaterally, without an impact on the post-session EEG. The degree
24 of enhancement in SMR was found positively related to the training course; the longer
25 patients took to attend for training the better was the SMR amplitude gain. Outcome benefits
26 included fewer awakenings and a longer time spent in slow wave sleep. However, the
27 majority of findings were correlational relating to individual differences in the SMR training
28 gain which correlated positively both with the increase in spindle frequency (13 – 15 Hz, C3)
29 in NREM sleep and with over-night memory consolidation, see Figure 5, which in turn
30 correlated positively with the increased frequency of spindles. Importantly the
31 counterbalanced design allowed within-subject comparisons of the impact of SMR versus
32 sham training in groups differentiated by condition order. For both there was evidence that
33 sham training had deleterious effects on the number of awakenings and of the time spent in
34 slow wave sleep.
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Figure 5 about here.

2.2. Alpha Activity.

Since the original EEG-biofeedback studies viewing alpha synchronisation as an index of relaxation, while desynchronisation remains associated with excitation, contemporary neuroscience has regarded alpha synchronisation as an active top-down inhibitory process for the exclusion of conflicting or irrelevant input (von Stein et al, 2000; Cooper et al, 2003; Basar, 2006) rather than as a brain idling index. The neural mechanisms underlying alpha production are incompletely understood with the classical view of thalamo-cortical circuits giving way to distributed alpha networks (Basar et al, 1997; Lopes da Siva et al, 1997; Klimesch, 1999). Different psychological associations have been posited between upper and lower alpha (Klimesch et al, 1993,1996; Petsch et al, 1997). Furthermore peak alpha frequency has been positively associated with intelligence and negatively with ageing (Jausovec et al, 2001; Clarke et al, 2004). Indeed the considerable individual differences in the alpha peak have led to an individual adjusted frequency procedure with alpha and sometimes adjacent frequencies (IAF; Klimesch et al, 1993). These are amongst the factors that have informed new protocols for neurofeedback, and see Bazanova and Vernon (2013, this issue) for further aspects of alpha methodology.

2.2.1. Visuospatial Rotation: High Alpha.

Informed by a transmagnetic stimulation (TMS) study in which mental rotation of a cube was enhanced by the TMS frequency which had been determined by the individual's alpha peak frequency (Klimesch et al, 1993), and secondly by the evidence that good cognitive performance was associated with high resting alpha power/low task-related alpha power and with the opposite relations in theta power (Klimesch, 1999), Hanslmayr et al (2005) explored raising high-alpha power versus reducing theta power (F3, Fz , F4 and P3, Pz, P4) both individually adjusted, in a counterbalanced design with outcome assessed

through spatial rotation. In a single-session experiment subjects ($N = 18$) were trained in separate 20-minute blocks of trials separated by the cognitive task. Results were analysed for the nine subjects with successful alpha learning and the ten with theta learning; only four in the alpha group were in the theta group. Raising upper alpha improved spatial rotation accuracy by about 10% post alpha training, and importantly there was a positive correlation between them ($r = 0.41 - 0.55$, $p < 0.05$), while learned control of theta reduction was not related to the behavioural or EEG outcome. EEG monitoring during the post-training assessment disclosed that the increase in upper-alpha power was sustained up until task onset but bore no relation to phasic alpha power induced by the task, in line with an influence of upper-alpha neurofeedback training on tonic rather than phasic EEG power. An increase in alpha at the right parieto-occipital placements, a primary locus for spatial rotation, correlated with the gain in upper-alpha power, offering the possibility that had this region been used for feedback larger improvements in mental rotation might have been obtained.

Figure 6 about here.

Zoefel et al (2011) went on to compare in five daily sessions visuospatial rotation in subjects ($N = 14$) trained to raise their upper-alpha amplitude versus a non-training control group ($N = 10$) with the alpha band individually adjusted each day. Successful training with feedback from bilateral parieto-occipital electrodes (P3, Pz, P4, O1, O2) was observed in 11/14 subjects, such that there was a linear increase in amplitude within/across sessions, and the baseline was higher in the fifth session than in the first session, demonstrating cumulative effects of learning in the tonic EEG that carried over from day to day. In support of training-bandwidth specificity the increase in the upper alpha band was independent of adjacent alpha and beta bands. Rotation was elevated only with neurofeedback and training also led to a higher upper-alpha amplitude than in controls.

2.2.2. Upper-alpha synchronisation and memory.

Upper-alpha training has also been explored for effects on working memory with an adaptation of the conceptual span test of Haarmann et al (2003). Escolano et al (2011) trained participants on five consecutive days with five 15-minute blocks and feedback averaged from parietal and occipital placements. IAF was computed daily from the 7.5 – 12.5 Hz band while changes in saturation of a fixation square were counted (active task) and comparisons were made between the 6/9 who were capable of increasing their upper alpha band (IAF + 2) and six no-training controls. Working memory was improved following training. Though within-session changes were not significant there was clear evidence of feedback learning across all twenty-five session blocks, and there was evidence of learning carrying over from day to day when comparing the first and fifth sessions during both the pre-session active task and a passive task without counting. In comparing whether there was training band specificity by examining lower alpha (IAF -1 to -3) and low beta (IAF +3 to +5), while independence was found for the active task, in the passive task there was a spread to include lower alpha.

Working memory has also been examined with digits backwards and forwards as an outcome measure (Nan et al, 2012) comparing sixteen participants who received upper-alpha Cz training based on IAF with sixteen non-training controls. They were given short condensed trials where in each session there were ten twenty second trials separated by five seconds when cognitive strategies were written down. There were three or four sessions within a day and a total of twenty sessions within fifteen days. In addition to the memory task pre and post-assessments consisted of eyes open and closed resting EEG (0.5 – 16 Hz). The relative amplitude increase in all bands between theta and sigma was positively correlated with session number while there was a negative correlation with delta. Memory improved with training and the improvement correlated positively with the increase in relative alpha power in the trained band only, in support of training-band specificity. There was no change

1 in the post-training resting EEG. Mental strategies were mostly positive such as thoughts of
2 love, family and friends.
3

4 2.2.3. Alpha Desynchronisation and Procedural Learning. 5

6 The impact on procedural learning of training down the mu rhythm (8 – 12 Hz) from
7 the right motor cortex (C4) in a single 30-minute session was examined in ten dextral
8 subjects by Ros et al (2013). The experiment was designed as an adjunct to a TMS
9 neurofeedback experiment (Ros et al, 2010) with a TMS protocol shown to increase
10 corticospinal motor evoked potentials and reduce short-interval intracortical inhibition (see
11 Gruzelier, 2013b). Serial left hand reaction-times were obtained with a task developed to
12 assess implicit procedural memory through repeating sequences intended to be unrecognized
13 by the participants (Nissen and Bullemer, 1987). The task was performed twice in a
14 counterbalanced design separated by a day, and on one occasion was preceded by
15 neurofeedback. Reduced RTs followed training, suggesting that alpha desynchronisation
16 acted to prime the motor cortex facilitating procedural learning.
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33

34 2.3. Feature Binding and Fluid Intelligence: Gamma and Beta Training. 35

36 Keizer et al (2010a,b) set out to revisit earlier evidence (Bird et al, 1978) of the ability
37 to modulate gamma activity with neurofeedback, an ability found to be sustained over 2 – 3
38 years (Ford et al, 1980). With outcome measures of feature binding (Hommel, 1998) and
39 fluid intelligence (Raven, 1938) Keizer et al (2010a) compared a group (N = 7) trained over
40 eight sessions to increase gamma (36 – 44 Hz) without increasing beta (12 – 20 Hz) with a
41 group (N = 7) trained to do the opposite - increase beta without increasing gamma. Occipital
42 midline electrode (Oz) training was obtained with testing spread across 10 to 11 days.
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Elevation of gamma with a maximum between 30 – 60 Hz was successful for the
experimental group reaching a plateau on the 6th session, with a positive correlation between
increases in power and in intelligence scores ($r = 0.82$, $p < 0.05$), a relation which extended

from 16 – 60 Hz, much wider than the gamma training-up band and including the beta inhibit band, see Figure 7. However, inhibition was found below the inhibited band where there was a reduction in slower frequencies below 10 Hz and importantly this reduction disclosed a suggestive negative correlation with slow wave 2-4 Hz power ($r = -0.70$, $p < 0.08$). While the comparison group was unsuccessful at increasing beta (12 – 20 Hz) and reducing gamma, there was an increase in slower wave activity maximal at 10-12 Hz which was in the opposite direction to the experimental group. Compatible with this there was a group difference in the RT episodic binding measure; this was reduced in the experimental group (mean 14.6 v 10.4) while doubling in length in the comparison group (mean 13.5 v 26.3).

Figure 7 about here.

Keizer and colleagues (2010b) went on to take into account frontal (Fz) as well as visual (Oz) functions given that the gamma influenced executive control processes are frontally situated. Coherence between the cortices was examined following the implication that fast wave activity in the gamma range acts focally, whereas slower activity acts over longer distances (von Stein et al, 1999; Varela et al, 2001). One important procedural modification involved restricting training to the primary band of interest, rather than requiring simultaneous control of gamma and beta as in the earlier experiment. Furthermore the two procedures differed in that the gamma feedback was from the occipital electrode only while beta feedback was the average of the two sites, occipital and frontal. Fifteen subjects received between 7 and 8 sessions of training, with pre and post-assessments as before and a recognition memory task adapted from Cycowicz et al (2001) which distinguished between familiarity and recollection. This task had provided evidence, albeit equivocal, for a possible double dissociation between the two EEG bands and between knowing and recollection processes in long-term memory (c.f Burgess and Ali, 2003; Gruber et al, 2008).

While the full bandwidth spectrum in this experiment was not disclosed, a lack of learning specificity vis a vis band and topography was disclosed by comparing the first and last sessions at the two sites. Oz gamma training evinced learning at the Oz training site across sessions but only within the last session, and Oz training influenced the frontal Fz non-training site where there was across-session learning in gamma and within-session learning in beta1, disclosing a spread of influence beyond the gamma training band. In contrast beta1 training, where feedback was the average of both frontal and occipital derivations, resulted in within-session learning frontally, but despite this there was no evidence of learning across sessions at either Oz or Fz. Possible interactions between sites were examined with coherence analysis. Within-session gamma coherence increased in session one for both groups, but this was not seen in the last session. On the other hand across sessions gamma Oz-Fz coherence increased in both groups, while beta coherence was increased only by beta training. Turning to the cognitive outcome the binding results of the previous experiment were replicated insofar as shape-location and location-response integration showed reduced costs following gamma feedback training, moderating the effects of task-irrelevant feature bindings, while the recognition memory task results disclosed the hypothesised double dissociation with gamma training improving recollection and beta training familiarity.

2.4. Theta (eyes-open).

While classically theta was associated with drowsiness, there has been more recent evidence of theta's role in learning and memorial processes and effort in attention. Theta facilitates long-term potentiation (Pavrides et al., 1988), is involved in recognition memory in encoding and recognition (Burgess and Gruzelier, 1997; Klimesch et al, 1997), is related to working memory load (Grunwald et al, 2001) and has been invoked in the encoding and retrieval of information in working memory (Klimesch, 1996,1999) and in linking a parietal-prefrontal circuit both for storage and up-dating (Sarnthein et al., 1998; von Stein et al., 1999;

von Stein and Sarnthein, 2000). Theta activity has also been implicated in memory consolidation through sleep (Buzsaki, 2004; Mitchell and McNaughton, 2008), Walker, 2004; Rauch, 2005). Frontal theta, while associated with increasing memory load, also synchronises with task demands, focussed attention, sustained concentration and meditation (Inouye et al., 1994; Grunwald et al., 2001; Jensen and Tesche, 2002; Onton et al. 2005; Kubota et al., 2001; Missonnier et al. 2006; Lagopoulos et al., 2009).

2.4.1. Frontal Theta.

In a methodological experiment Enriquez-Geppert et al (2013) set out to explore the trainability of anterior theta in thirty one participants who were randomised to either training-up theta or to a pseudo feedback control group, following implications that frontal theta recorded from the midline has potential as a marker of executive functions (Mitchel et al, 2008) including cognitive control (Cavanagh et al, 2011). With no cognitive/affective outcome measures this will be outlined in Part III (Gruzelier, 2013b), however, an application of training up anterior theta with the elderly (Wang and Hsieh, 2013) is reviewed in section 4.2.

2.4.2. Central (Cz) Theta.

As indicated above (2.1) Vernon et al (2003) alongside SMR training also examined eight sessions of eyes open Cz theta up-training while inhibiting both delta (0-4 Hz) and alpha (8 – 12 Hz), along with a non-training control group. Theta training provided no gains for a two-digit CPT or a semantic working memory task whereas SMR training was beneficial, while neither neurofeedback protocol facilitated performance on a three-digit CPT with the greater memory load. Whereas there was evidence of learning with SMR training this was not the case with training-up theta, in fact theta was effectively trained down for within sessions the desired theta/alpha ratio went significantly in the opposite direction with alpha increasing and theta decreasing, not vice versa, and without improvement in the

1 theta/delta ratio. Most likely the broad band alpha inhibit was inappropriate given the
2 involvement of alpha in a diversity of task related processes (Klimesch et al, 1998;
3
4 Doppelmayr et al, 2002) that would encroach on neurofeedback learning.
5
6

7 2.4.3. Posterior Theta. 8 9

10 Classical associations of theta in relation to the arousal spectrum have involved
11 posterior recordings with posterior theta indexing low arousal, tiredness and inattention.
12 Beatty and colleagues (1974) had reported that individuals receiving theta (4-7 Hz) up-
13 training versus down-training (O1, P3), decreased and increased respectively their radar
14 detection performance. Since then posterior theta activity has been implicated inter alia in
15 consolidation of memory processes through sleep (Buzsaki, 2002; Mitchell and McNaughton,
16 2008), Walker, 2005; Rauch et al, 2005). Reiner et al (2013) explored the impact of a single
17 session of training up eyes-open theta on the learning of a 30-minute finger-thumb apposition
18 task and its repetition following one, two, three and seven nights of sleep. Participants (N =
19 38) were randomised to 30 minutes of Pz theta (4-7Hz) or beta 1(15 – 22Hz) training or to a
20 control group who watched movies. Speed of performance improved in all groups with
21 practice, but more so following theta training. There was an immediate advantage of (8.7%)
22 which was sustained and saw an additional 8.5% gain between the last two sessions
23 compared with a 4% gain in controls and no improvement in the beta group. Whereas the
24 initial gains were attributed to synaptic consolidation, the final delayed gain was
25 hypothesised to reflect consolidation at a systems level (Diekelmann and Born, 2007) with
26 enhanced performance through the redistribution of new memories to multiple, distributed
27 brain regions (Song, 2008).
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52

53 That theta training was central to the improvement in speed of performance (there were
54 no findings with accuracy) was supported by positive correlations between the ratio of theta
55 to beta, a ratio obtained to normalise for individual differences in absolute theta, and the
56
57
58
59
60
61
62
63
64
65

performance gain on all assessments following the first night's sleep, but not immediately after training. Taking the results together it was inferred that consolidation of learning during sleep was responsible for the gains following theta training, though a control for practice will be necessary to validate the involvement of sleep in the process. However, this does not detract from the main finding that theta neurofeedback contributed to the improvement in motor procedural learning.

3. Mood.

3.1. Alpha/Theta Training.

In establishing what was the first evidence of operant control over the theta/alpha ratio with eyes closed, Egner et al (2002) had shown in healthy subjects that when comparing a contingent with a non-contingent control A/T protocol, both protocols led to identical ratings on Activation Deactivation self-report scales (Thayer, 1967), whereas only contingent training produced an increase in the theta/alpha ratio. One implication from this was that there was likely to be more to the reported clinical and performance benefits of A/T training than relaxation per se. Indeed performing arts studies with adult musicians, dancers, and actors, and child musicians, which will be reviewed in Part II (Gruzelier, 2013a, and see Gruzelier, 2012), have shown not only enhancements in Creativity and Technique, but also enhancements with a strong emotional component reflected in Communication in performance which was underpinned by ratings of Enjoyment, Confidence, and Emotional Commitment and Conviction (Egner and Gruzelier, 2003; Gruzelier et al, 2013a,b; Gruzelier, 2012).

Raymond et al (2005) went on to examine possible benefits for well-being in socially anxious and withdrawn students randomised to alpha/theta training or to a mock training control where feedback was non-contingent and outcome assessments were examined with

the Profile of Mood States (McNair et al, 1992). Within-session learning but not between sessions was highly significant and was combined with advantages favouring contingent A/T feedback on self-ratings of Composure, Agreeableness, Elevated Mood, Confidence and Energy. The control group, who on debriefing were unaware of their control status and whose perception of learned control was on a par with the contingent group, showed no evidence of learning and reported more Composure and Tiredness. A/T training produced a 25% improvement in mood overall.

3.2. SMR Ratio Training.

SMR enhancement has also been shown to have an impact on mood (Gruzelier, 2013c). In the first attention/ERP study with music students (Egner and Gruzelier, 2001; section 2.1) an activation/ deactivation checklist (Thayer, 1967) was examined at the beginning and end of each of the SMR and beta1 protocols which were administered for 15 minutes in a counterbalanced order in the same session for a total of 10 training sessions. Whereas with both protocols Tiredness was increased within sessions, Calmness was differentially affected with an improvement following SMR training only. This was theoretically consistent with the calming influence of the trained putative inhibitory influences of the SMR rhythm on the sensorymotor cortex with the consequence of reduced motor interference and reduced muscle tone (Serman, 1996), and was endorsed by the post-training reduction in the musician's impulsivity indexed by commission errors in a continuous performance test (Egner and Gruzelier, 2001). The increase in calmness was also consistent with an experiential analysis conducted with some of the musicians from a taped semi-structured interview (Edge and Lancaster, 2004). Five out of the ten musicians reported discernible experiences about the SMR/beta1 training which the authors encapsulated by a report of one student that it 'made the mind breathe'.

Further evidence of beneficial influences of SMR neurofeedback on mood was reported by Ros et al (2009) in the microsurgery study with benefits for both anxiety and surgical skills. In those receiving A/T training the increase in the theta/alpha ratio, which correlated with an efficient performance, also correlated with a reduction in anxiety. Anxiety has been similarly reduced in twenty female swimmers described as adolescents with 5-6 years of professional experience (Faridnia et al, 2012) as assessed by a Sport Competition Anxiety Test (Martens, 1997) which was administered before and after neurofeedback or a control condition the details of which were not reported. Here twelve sessions of SMR (12-15 Hz) ratio training were combined with beta1 (15- 18 Hz) training, both protocols with theta (4-8 Hz) and beta2 (22-37 Hz) inhibits. They were given in two 20-minute blocks within a 45 minute session with order of protocols not reported. Reduced anxiety scores were found solely in the neurofeedback group. Given differential influences of SMR and beta1 ratio training on mood (Gruzelier, 2013c), separation of SMR from beta1 training may have been informative.

3.3. Alpha Synchronisation.

Alpha training (8 – 12 Hz) was revisited by van Boxtel and colleagues (van Boxtel et al, 2012) developing a user-friendly ‘relaxation’ device suitable for use in public with wireless electrodes mounted in an audio headset placed over the middle of the head allowing recording approximately from the central electrode chain. An alpha group (N = 18) was compared with a random beta group (N = 12) who heard a 4 Hz band between 14 and 30 Hz which differed each day, and a control group who listened only to music (N = 20). In each session the mostly college student participants listened to their choice of music which was only heard clearly when the training frequency band was enhanced. Music ranged over hard rock, easy listening and classical, assumed to have identical effects on relaxation, but with no empirical verification or matching of groups for choices. There were fifteen sessions within

1 four weeks, and participants were trained in groups of five. The participants were instructed
2 to 'sit back and relax' with their eyes open. Training sessions were divided into three blocks
3
4 of 8 minutes bracketed by 5 minutes of one of four cognitive tasks in the fixed order of
5
6 flanker task, stop-signal task, Stroop task and N-back task, and interspersed with ten
7
8 subjective relaxation ratings. Roughly a week before and after training and again at an
9
10 approximately four month follow-up, a topographical EEG was recorded with eyes open and
11
12 closed. In addition questionnaires were completed encompassing mood, sleep and quality of
13
14 life.
15
16
17
18

19 Importantly the recording of alpha from the headset did not differ from the laboratory
20
21 recording from C3/C4 derivations. There was no statistical evidence of relaxation comparing
22
23 session 1 and 15 on a rating scale, nor group differences, and in an exit interview while about
24
25 half (53%) in the alpha group mentioned that the training was relaxing which was twice the
26
27 number in the other groups this was not statistically significant, nor were there group
28
29 advantages in questionnaire measures of mood, sleep and quality of life. Relative alpha
30
31 power was averaged across lateral central, parietal and occipital placements in eyes open and
32
33 closed conditions disclosing that parieto-occipital alpha was higher than central alpha. With
34
35 alpha training eyes-closed alpha increased more than in the beta group, but as beta training
36
37 would be activating this would be expected, while with eyes open the training course had an
38
39 identical effect in the alpha and beta groups in raising alpha, which was increased further by
40
41 follow-up more so in the alpha group. Interpretation of the results is hampered by an absence
42
43 overall of paired comparisons within and between groups, and inspection of the figures
44
45 indicated that the music control group had considerably higher levels at baseline. Validation
46
47 of the effects of the individualised music choices on relaxation may lead to a less equivocal
48
49 outcome.
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

4. Applications.

4.1. Sporting Skills

Applications of neurofeedback to improve sporting performance and to compare elite with novice performance are at an early stage but will be apposite (Thompson et al, 2008). Little progress has been made since the pioneering studies of Landers et al (1991) who reported that the feedback of an EEG analysis of slow potentials improved performance in pre-elite archers. They followed evidence that in sportsmen prior to the execution of a skill there was a unilateral slowing of left temporal activity and randomised 24 pre-elite archers to three groups: a left temporal (T3) slowing experimental group, a right temporal (T4) slowing group controlling for expectancy and motivation, and no-training controls. Training involved a 45 -75 minute single neurofeedback session with shooting (27 shots) examined before and after. Differential outcomes occurred in the training groups with improved accuracy in the left temporal group, a deterioration in the right temporal group, and with no change in the no-training group while examining spectral power increases on the left did not differentiate the groups, whereas beta power increased on the right in the 13-30 Hz range in the right temporal group from which evidence of cognitive processing was inferred as seen in novices for elite performance becomes automatic (Fitts and Posner, 1967).

An individualised approach has been examined in six golfers in whom EEG (FPz) in conventional spectral bands was recorded immediately before golf putting (N = 80), contrasting successful with unsuccessful putting attempts in order to ascertain the optimal EEG profile associated with success that met with inter-rater agreement (Arns et al, 2008). Three daily training sessions were given consisting of four conditions of eighty putts each with feedback in the second and fourth conditions. In the feedback conditions a continuous tone was presented before each putt up until the optimal EEG pattern occurred - the signal to proceed with putting. Performance was better overall with feedback, but examination of the

three training sessions separately failed to show benefits in the last session; a preferential feedback effect occurred only on the second occasion in which there was evidence of cumulative learning and better performance overall. The conditions of play – outdoor or indoor – varied due to weather which may have contributed to the falloff in performance in the third, final training session. While some other reports are emerging, in the absence of controls, the validation of measures (e.g., Sherlin et al, (2013) and reporting of control procedures (e.g., Paul et al, 2012) little can be concluded.

4.2. Cognition in the Elderly.

A number of exploratory attempts have been made for preserving cognitive functions in the healthy elderly, with the clear conclusion that age does not exclude neurofeedback learning.

4.3.1. Alpha Training.

In an exploratory study of six participants aged 70 – 78 years Angelakis et al (2007) contrasted: i) increasing peak alpha frequency while inhibiting alpha amplitude (8 – 13 Hz) which was successful in 2/3, ii) increasing alpha amplitude while inhibiting peak frequency, successful in 2/2, and iii) one mock feedback control. Thirty one to thirty five sessions were given once or twice a week involving a total of 24 minutes of posterior training (POz). There were impressions of differential effects on outcome measures of processing speed, executive functions and memory, all known to decline with age, with an improvement from amplitude training with processing speed and executive function, and improvement from peak frequency training with visual, verbal and working memory. They noted that similar to Egner et al (2004), the outcome in the topographical EEG was expressed frontally and not at the posterior training site in the three with peak training, a result not seen in the other subjects.

1 Encouragingly it may be concluded that the age range 70 – 78 years is not a handicap to
2 regulating brain activity.
3
4

5
6 The implications of this pilot study were followed up by Lecomte and Juhel (2011)
7
8 with 30 participants aged between 65 and 85 years who were assigned to one of three groups:
9
10 an increase in upper-alpha (10 – 12 Hz) power while inhibiting theta (4 – 7 Hz; transverse
11
12 montage (C3-Cz/Cz-C4), essentially a relaxation control group performing simple yoga
13
14 movements, or a no-intervention control group. Sessions involved 30 minutes of training on
15
16 four occasions within a single week. Neurofeedback learning was on the whole successful
17
18 with a significant increase over sessions in alpha power seen in 8/10, together with an
19
20 increase in the alpha/theta ratio (7/10), without a significant reduction in theta (6/10). The
21
22 EEG was examined pre and post training along with delayed recall and semantic fluency and
23
24 a memory learning test, and following each session tension level was recorded with analogue
25
26 scales of stress and anxiety. Cognitive improvements occurred in all groups while tension
27
28 level fell in both training groups; none of the improvements were associated with
29
30 neurofeedback learning. The four training sessions were surely too few with elderly
31
32 participants.
33
34
35
36
37
38
39
40

41 4.3.2. Theta Training. 42 43 44

45 Following evidence that abnormally enhanced theta has been associated with the
46
47 greater cognitive impairments in patients with either mild cognitive impairment or dementia
48
49 (e.g. Babiloni et al, 2007). Becerra et al (2011) set out to reduce theta in fourteen normal
50
51 participants aged 60-85 years who had evidence of abnormally high theta in one lead.
52
53 Electrode placements were individualised according to the patient's maximum theta elevation
54
55 and involved the following locations for the sample: F4, C3, C4, P3, F7, F8, and T6. They
56
57 were randomly assigned to either an experimental group or a noncontingent feedback control
58
59
60
61
62
63
64
65

group. Thirty 30-min sessions were given over 10 – 12 weeks. In the experimental group theta power was successfully reduced and coincided with reductions in F3 and midline absolute theta power and left lateralised increases in alpha power. However, the control group also disclosed changes in relative theta and alpha power. Pre and post assessments consisted of the Wechsler Adult Intelligence Scale, neuropsychological scales of memory and executive functions, and topographical EEG. The neurofeedback group showed improved verbal IQ, executive functions and attention and shared memory improvements with the control group who also demonstrated a decrease in performance IQ. Links were made between normalising midline theta and improvements in attention, and between the reduction in theta at F3 and Fz and the increase in left-sided alpha regarding the improvement in verbal IQ, and the frontal changes for executive functions. Of relevance to the question of topographical specificity the reductions in theta seen in the post training EEG did not coincide with the training locations based on maximum theta power. The normalising of theta and the increase in alpha were interpreted as arising from thalamic influences (Hughes et al, 2004). That some benefits were shared with the control group, as found in the French study, indicate that aside from training, the elderly may be responsive to motivational and social factors through research engagement, while the inclusion of a sham control task ensured that processes involved in task engagement were also shared.

Wang and Hsieh (2013) examined Fz theta (4-7 Hz) training for its potential to improve attention and recognition memory in an older (61 to 72 yrs) community sample, described as high functioning elders engaged in community work and screened for dementia and depression, who were compared with university students (21 – 25 yrs). Subjects were allocated in groups of eight, to 12 sessions over four weeks of theta training or to random 4 Hz band training in the range 10 – 25 Hz, both protocols were with delta and gamma inhibits. Regarding characterisation, pre-training assessments indicated that the elderly were in

1 relatively good condition cognitively when compared with the students, with superior
2 Attention Network Task (Fan et al, 2002) accuracy scores, though RTs were slower (a
3 speed/accuracy trade off ?), recognition memory was poorer, while a topographical recording
4 of the resting eyes-open EEG showed less theta and delta power in the older group in the
5 fronto-central midline region (c.t. Beccera et al, 2011), which carried over to the post-training
6 assessment.
7

8
9 Importantly the elderly were capable of increasing midline frontal theta and learning
10 indices disclosed evidence of neurofeedback learning equally in both students and the older
11 sample, and in the EEG no difference between them at Fz; these effects were absent in the
12 control groups. Furthermore following neurofeedback measures showed improvements
13 specific to the elderly were found in orienting (RT ANT) and in recognition memory, while
14 ANT conflict was improved in both groups relative to the sham controls; there was no impact
15 on alerting, see Figure 7. Correlational data between the neurofeedback learning indices and
16 the cognitive outcome would have confirmed a meditational link with the theta training.
17
18

19 4.3.3. Gamma Training.

20
21 Further support for the ability of older subjects to engage with neurofeedback training
22 was obtained by Staufienbiel et al (2013) with twenty participants aged 58 – 80 years
23 allocated to either eight sessions of Fz gamma (36 – 44 Hz) or broad band low beta (12 – 20
24 Hz) training and with no more than two days between sessions. The main aim was to extend
25 to the elderly the improvement in fluid intelligence following gamma neurofeedback in
26 younger participants (Kiezer et al, 2010a), in view of gamma deficiency reports in the elderly
27 and in dementia (Werkle-Bergner et al, 2009; Stam et al, 2002), along with the differential
28 benefits from beta and gamma training respectively on familiarity and recollection in
29 recognition memory. Before and after the training course eyes-open EEG was examined
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1 across frontal, central and parietal chains and frontal and central chains within sessions,
2 together with a well-being questionnaire concerning appetite, need for sleep, concentration,
3 memory and mood.
4
5
6
7

8 There was evidence of within-session learning in both groups. Gamma training
9 increased both gamma and beta amplitude while beta training demonstrated band specificity
10 with a comprehensive increase in broad band beta. In contrast there was no evidence of
11 gamma learning across sessions and with only a tendency for beta amplitude to increase
12 following beta training (and to decrease with gamma training). In neither group was there an
13 impact on the post-training resting EEG, nor did gamma show specificity regarding other
14 outcome measures, whereas following beta training one notes a weak tendency for an average
15 27% increase in fluid intelligence, but one in keeping with Keizer et al's (2010a) correlation
16 between fluid intelligence and a broad 16 – 60 Hz band. Familiarity in recognition memory
17 increased in both groups supported by self-reported improvements in memory; a likely
18 reflection of engagement with the experimental procedure rather than a reflection of learned
19 control must be considered. The self-assessment questionnaire also disclosed in both groups
20 reduced appetite, and a tendency for a greater need for sleep following beta training, which
21 given the impact of SMR on sleep spindles may have arisen from inclusion of the SMR band
22 within the beta training band. Changes in gamma were not correlated with outcome, though
23 an increase in beta power at F4 was correlated with a reduction in appetite and an increase in
24 beta power at F4 and C4 correlated weakly with a smaller recollection memory score.
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52

53 5. Validation. 54

55 Having reviewed the efficacy of EEG-neurofeedback for optimising function in healthy
56 individuals, before concluding Part I there are two sources of evidence in support of
57
58
59
60
61
62
63
64
65

validation to be summarised. The first is where a neurofeedback experimental group is characterised by a uniquely successful cognitive and/or affective outcome when compared with a comparison group and/or a control group; should in the absence of a control group two neurofeedback protocols show the same benefits these are not included as the result may be a reflection of nonspecific influences on training. The second source is where there is evidence of neurofeedback learning coupled with affirmative inter-relations between learning and post-training assessments. This represents the stronger verification with as yet a smaller evidence base. Group differences favouring neurofeedback are ordered according to spectral training band, mindful that protocols may differ in methodological features such as electrode placements, inhibits and so on. Gains are assigned to the particular EEG band that discloses them rather than to the protocol training bands themselves should the latter not have been fruitful.

5.1. Cognitive and Affective Outcome.

5.1.1. SMR ratio.

From this the most extensively studied EEG-neurofeedback protocol, with clinical origins, a substantive range of cognitive and affective differential group advantages has been reported in controlled studies of healthy participants including:

- i) Sustained attention in children and adults along with enhancement of ERP indices of attention (Egner and Gruzelier, 2001, 2004; Vernon et al, 2003; Gruzelier et al, 2013a).
- ii) Memory in adults and children (Vernon et al, 2003; Barnea et al, 2005; Hoedlmoser et al, 2008).
- iii) Spatial rotation (Doppelmayr and Weber, 2011).
- iv) Reaction time and complex psychomotor skills (Doppelmayr and Weber, 2011; Ros et al, 2009).

v) Calmness and reduced performance anxiety (Ros et al, 2009; Faridnia et al, 2012; Gruzelier 2013c).

vi) Increased frequency of sleep spindles associated with improved memory (Berner et al, 2006; Hoedlmoser et al, 2008; Schabus et al, 2013).

The various outcomes with healthy participants provide support for SMR training in the treatment of ADHD for information processing, affect regulation and sleep (Monastra et al, 2005; Arns and Keneman, 2013). There are also promising applications in the performing arts, especially for novice performance requiring lower-level skills when compared with elite performance, which will be outlined in part II (Gruzelier, 2013a).

5.1.2. Beta and Gamma.

Surprisingly, given that the classical EEG response to cognitive and affective engagement is the production of beta activity through desynchronisation of the EEG (Ray and Coles, 1985), and that up-training of the 'beta'/theta ratio is a standard ADHD protocol, there have been few investigations of beta training to date with healthy participants, or it has been of secondary interest, or included as a control protocol.

Beta together with gamma training has resulted in a number of benefits:

- i) Egner and Gruzelier (2001) found that beta1 shared with SMR learning positive relations with an increase in the P300b amplitude, a gain which in the case of beta1, but not SMR training, was replicable (Egner and Gruzelier, 2004) and compatible with enhanced arousal following beta1 training.
- ii) A comparison of gamma (36 – 44 Hz) and broad low beta band (12 – 20 Hz) protocols disclosed improved recollection in long-term memory and perceptual binding following gamma training and improved familiarity in

1 recognition memory following beta training (Keizer et al, 2010b), with the
2 proviso that training-up gamma also increased beta (12 – 20 Hz) amplitude.
3

- 4
5 iii) In the elderly there was a tendency for a gain in fluid intelligence following
6
7 broad-band beta training (Staufenbiel et al, 2013), in support of Keizer et al
8
9 (2010a), Figure 7.
10

11
12 In view of the gains following beta training caution is warranted when beta training is
13
14 used for secondary purposes such as a control for nonspecific influences on training.
15
16 Similarly the use of random narrow band beta as a control procedure (van Boxtel et al, 2012)
17
18 must be applied with caution when considered, firstly with the extensive evidence of one-trial
19
20 neurofeedback learning (Landers et al, 1991; Hanslmayr et al, 2005; Berner et al, 2006; Ros
21
22 et al 2010, 2012, 2013; Reiner et al 2013), and secondly with the leakage of training band
23
24 influences within the beta spectrum (Keizer et al 2010a,b). Furthermore given relations
25
26 between arousal and beta, over-arousal may result and reduce performance in line with the
27
28 Yerkes Dobson inverted-U relation between arousal and performance (Duffy, 1957;
29
30 Anderson, 1990).
31
32
33
34
35

36 5.1.3. Theta.

37
38 Theta protocols have provided promising outcomes:
39
40

- 41 i) Training-down theta with placements individualised on the basis of EEG theta
42
43 maxima coincided with improvements in verbal IQ, executive functions and
44
45 attention in the elderly (Becerra et al, 2011).
46
47
48 ii) Training-up frontal theta in university students and a healthy elderly sample
49
50 improved executive attention in both groups and improved recognition memory
51
52 and orienting in the elderly (Wang and Hsieh, 2013).
53
54
55 iii) One session of training-up theta at Pz improved motor procedural learning with
56
57 additional gains following sleep (Reiner et al, 2013).
58
59
60
61
62
63
64
65

- iv) Training-up the theta/alpha ratio with eyes closed with children led to a reduction in impulsive errors in a continuous performance test (Gruzelier et al 2013a).
- v) Training-up the theta/alpha ratio with eyes closed led to improvements in mood and well-being in students (Raymond et al, 2005a).

In terms of affective processes wide ranging benefits encompassing ratings of Composure, Agreeableness, Elevated Mood, Confidence and Energy on the profile of Mood States (McNair etc), provide support for earlier clinical studies where the A/T protocol was incorporated with other treatment procedures (Peniston and Kulkowsky, 1989, 1990, 1991; Saxby and Peniston, 1995). The influence of A/T training on mood would appear to go beyond anxiety reduction and raise affect and energy levels, boosting self confidence and commitment in performance, as well as psychological integration (Raymond et al, 2005; Gruzelier, 2009, 2013a). Interestingly superior A/T learning has also led to cognitive gains including improved surgical performance aside from a reduction in performance anxiety, and substantive improvements in sustained attention in children (Gruzelier et al, 2013). But by far the more extensive evidence is the impact on creativity in the performing arts which is outlined in Part II (Gruzelier, 2013a).

5.1.4. Alpha Power.

Successful outcomes have included:

- i) Gains in spatial rotation through the training-up of the combined frontal/parietal chains (Hanslmayr et al, 2005) and through posterior alpha training (Zoefel et al, 2011).
- ii) Working memory facilitation by increasing posterior upper-alpha power (Escolano et al, 2011), and with a centrally located derivation (Nan et al, 2012).

iv) In contrast training down broadband alpha in the form of the mu rhythm from motor cortex producing EEG desynchronisation facilitated implicit procedural memory indexed by reduced RTs compatible with increased arousal (Ros et al, 2013).

Training upper-alpha power holds promise for further applications. The question has been posed (Hanslmayr et al, 2006) as to whether upper-alpha enhancement may be responsible for gains following SMR training due to upper-alpha falling within the SMR band, given the individual differences in the peak alpha frequency, which tend not to be taken into consideration with the IAF procedure when other bands have been trained. However, firstly in the upper-alpha training studies involving parietal/occipital training Nan et al (2012) reported a broad-band increase extending from theta to sigma whereas it was only the upper-alpha band that correlated with improved memory, while Zoefel et al (2012) reported independence of upper-alpha from the sensory-motor rhythm band. Secondly the induced cognitive effects of SMR training (2.1) would be much broader than the contemporary views of upper-alpha correlates (Klimesch, 1999). Nevertheless studies contrasting upper-alpha and SMR training and also with lower-alpha will be informative.

5.2 Relations between Neurofeedback Learning and Outcome.

In this penultimate section of Part I the focus will be on the evidence of neurofeedback learning indices that correlated with improvements in outcome assessments. This demonstration is of singular significance because the correlation between a learning index and improvement in an outcome assessment provides support for the crucial mediation link between performance enhancement and feedback learning.

There are seven studies reporting affirmative correlations between learning and outcome measures out of the twenty-three controlled studies that have reported group

1 advantages in favour of neurofeedback coupled with evidence of neurofeedback learning. In
2 addition to the seven reports there are others which pertain to creativity and the performing
3 arts which are reviewed in Part II. Learning assessment will be taken at face value and no
4 distinction will be made between within-session learning and cumulative learning across
5 sessions, issues that will be considered in detail in Part III. Here the seven studies are
6
7 categorised by neurofeedback protocol and they provide nine instances in all of feedback
8
9 learning-outcome relations (numbered i – ix).
10
11
12
13
14
15

16 SMR ratio training has been correlated with outcome in three of the seven studies
17
18 (Egner and Gruzelier, 2001; Ros et al, 2009; Schabus et al, 2013).
19
20

- 21 i) In the earliest report (Egner and Gruzelier, 2001) a positive impact was seen in
22 a reduction of impulsive errors ($r = -0.79$, $p < 0.001$; see Figure 1; 2.1.1.)
23
24 contributing to an enhanced d-prime in a continuous performance task, and in
25
26 an enhanced P300b ($r = 0.49$, $p < 0.06$), see Figure 2.
27
28
- 29 ii) The reduction in impulsive errors was theoretically consistent with a
30
31 subsequent study showing correlations between SMR learning and outcome
32
33 (Ros et al, 2009). Ros et al (2009) with junior doctors found a correlation
34
35 between SMR learning and the pause time in between microsurgical tasks in a
36
37 simulated cataract operation ($r = 0.72$, $p < 0.019$; 2.1.2). This was in keeping
38
39 with a more modulated performance, and was associated with more efficient
40
41 surgical skills following SMR training.
42
43
- 44 iii) The third source of SMR correlational evidence was from a sleep study
45
46 (Schabus et al, 2013) showing inter-correlations between SMR learning and
47
48 memory improvement over night ($r = 0.40$, $p < 0.027$; Figure 5a,b), and
49
50 increased sleep spindle frequency (13-15 Hz) in NREM sleep ($r = 0.38$, $p <$
51
52 0.038). These relations supported the sensory-motor rhythm affinity with
53
54
55
56
57
58
59
60
61
62
63
64
65

sleep spindles and the role of sleep in memory consolidation (Niedermeyer, 2005; Shabus et al, 2004).

- iv) Beta1 enhancement correlated with the amplitude of the P300b in auditory detection ($r = 0.55$, $p < 0.05$, see Figure 2, (2.1.1.), in support of increased arousal (Egner and Gruzelier, 2001).
- v) When training gamma (36 – 44 Hz) without increasing ‘beta’ (12 – 20 Hz) Keizer et al (2010a) reported a correlation ($r = 0.82$, $p < 0.05$) between fluid intelligence and a broad beta/gamma band (16 – 60 Hz), see Figure 7 (2.3).

Upper-alpha has demonstrated mediation links in two studies.

- vi) Hanslmayr et al (2006) reported that one session of training to increase upper-alpha and decrease theta led to improvement in spatial rotation which correlated with the gain in upper-alpha ($r = 0.41 - 0.55$, $p < 0.05$; see Figure 6).
- vii) Nan et al (2012) reported that upper-alpha training correlated with gains in a working memory task: digits forwards ($r = 0.501$, $p < 0.05$), digits backwards ($r = 0.543$, $p < 0.05$).

Finally theta up-training at Pz, both with eyes closed and with eyes open has been correlated with cognitive and affective outcome.

- viii) Training the theta/alpha ratio with eyes closed, Ros et al (2009) in the micro-surgical skills study reported that elevation of the ratio was positively associated both with the improvement ($r = 0.64$, $p < 0.047$) and overall efficiency in technical psychomotor skills ($r = -0.52$, $p < 0.06$) as well as with a reduction in performance anxiety ($r = -0.66$, $p < 0.053$). Here the group who elevated the theta/alpha ratio shared the post-training outcome benefits with

the group receiving SMR ratio learning; the latter incidentally included down-training theta with eyes open.

- ix) Reiner et al (2013) found that training-up posterior theta with eyes open correlated with gains in procedural learning following one, two, and six nights of sleep ($r = 0.64 - 0.66$, $p < 0.01$) but not immediately after training (0.33, ns). From this it was inferred that the theta synchronisation protocol facilitated memory consolidation processes in sleep. Benefits for memory consolidation following sleep have also been shared with SMR learning as found by Schabus et al (2013), in this case an SMR protocol without theta inhibits in contrast to the conventional SMR ratio protocol which shared benefits with theta training (Ros et al, 2009). The parallels between the contrasting theta protocols and SMR training are not understood.

One cannot necessarily infer from an absence of reportage an absence of learning because there has not been a tradition of considering this. There is also the proviso that because the understanding of the nature of neurofeedback learning is by no means a clear cut, well trodden issue, it is at an early stage and learning may have eluded measurement. For example, learning as in a carry-over to successive pre-training baselines may have gone unnoticed (Gruzelier, 2013b, Part III). The failure to find learning may be a reasonable inference when no cognitive/affective advantage has been disclosed as in the up-training of theta in the report of Vernon et al (2003). Additionally attribution of learning to the neurofeedback process per se may not always be the case, when to give an example, learning falls away within a session once training stops, and there is no incremental increase across sessions either during training or in successive pre-training baselines. This example may reflect a process incidental to learning such as concentration or effort in attention in the feedback process.

While the number of studies is small, such evidence is a methodological advance for the neurofeedback field in general, one which has been seldom encountered in the more populous clinical literature.

5.3. Validation Conclusion.

As has been documented in this review successful validation within this re-emergent EEG-biofeedback domain has included diverse protocols: SMR and SMR ratio, beta1 and beta1 ratio, alpha/theta ratio with eyes closed, upper-alpha, alpha desynchronisation, gamma and gamma ratio, frontal theta up-training, posterior theta up-training and down-training theta EEG maxima. Approaches have arisen from contemporary neuroscience offering new opportunities in the clinic and the field. Validation has also included standard clinical protocols extending their repertoire and enriching their clinical potential, as exemplified by the feasibility of conducting neurofeedback with the elderly and the multiplicity of approaches applied: increasing high alpha power while inhibiting an increase in theta (Lecomte and Juhel, 2011), training-down individualised theta maxima (Becerra et al, 2011), training-up frontal midline theta with delta and gamma inhibits (Wang and Hsieh, 2013), training-up frontal gamma (Staufenbiel et al, 2013), and increasing alpha power while maintaining peak alpha frequency or vice versa (Angelakis et al, 2007). Their impact on cognitive and affective outcome in controlled studies with the elderly has included: executive functions, IQ and attention (Becerra et al, 2011) and recognition memory, orienting and executive attention control (Wang and Hsieh, 2013).

These laboratory and field studies with healthy volunteers have raised a host of methodological and theoretical issues that will stimulate research and assist in understanding the mechanisms underlying neurofeedback learning and the nature of the EEG and its cognitive/affective relations, which in turn will refine protocol implementation and real world applications. In Part III such issues will be enumerated further including inter alia the

1 independence of training bands, behavioural outcome and topography, experimental design
2 and issues surrounding what may be regarded as the elephant in the room – the fact that
3
4 neurofeedback is a learning process that must be guided and nurtured.
5
6

7 In summary, the broad encompass of cognitive and affective outcome assessments from
8
9 controlled studies has favoured:
10

11 sustained attention, orienting and executive attention, the P300b, memory, spatial
12
13 rotation, RT, complex psychomotor skills, implicit procedural memory, recognition memory,
14
15 perceptual binding, intelligence and wide ranging aspects of mood and well-being.
16
17

18 The mediation links through correlations between neurofeedback learning and outcome
19
20 gains have included:
21
22

23 sustained attention, P300b, working memory, mental rotation, motor procedural
24
25 learning, psychomotor skills, fluid intelligence and anxiety in performance.
26
27

28 There is now sufficient evidence validating the role of EEG-neurofeedback in
29
30 enhancing function to dispel the lingering vestige of prejudice against the value of this EEG
31
32 methodology. Meanwhile, neurofeedback for optimising function has materialised centre-
33
34 stage in neuroscience through the advent of fMRI applications (Ruiz et al, 2013).
35
36
37
38
39
40

41 References.

42
43 Anderson, K.J. (1990) Arousal and the inverted-U hypothesis: A critique of Neiss's
44
45 "Reconceptualizing arousal." Psychological Bulletin, 107, 96-100. doi: 10.1037/0033-
46
47 2909.107.1.96
48
49
50
51 Angelakis, E., Lubar, J.F., Stathopoulou, S., 2004. Electroencephalographic peak
52
53 alpha frequency correlates of cognitive traits. Neuroscience Letters, 371, 60-63.
54
55
56
57
58
59
60
61
62
63
64
65

Arns M., Kleinnijenhuis, M., Fallahpour, K., Breteler, R. (2008) Golf performance enhancement and real-life neurofeedback training using personalized event-locked EEG profiles, *Journal of Neurotherapy*, 11, 11-18.

Arns, M., de Ridder, S., Strehl, U., Breteler, M., Coenen, A. (2009) Effects of neurofeedback treatment on ADHD: the effect on inattention, impulsivity and hyperactivity: a meta-analysis. *Clinical EEG Neuroscience*, 40, 180-189.

Babiloni, C., Squitti, R., Del Percio, C., Cassetta, E., Ventriglia, M.C., Ferreri, F., Tombini, M., Frisoni, G., Binetti, G., Gurzi, M., Salinari, S., Zappasodi, F., Rossini, P.M. (2007) Free copper and resting temporal EEG rhythms correlate across healthy, mild cognitive impairment, and Alzheimer's disease subjects. *Clinical Neurophysiology*, 118, 1244-1260.

Barnea, A., Rassis, A., Zaidel, E. (2005) Effect of neurofeedback on hemispheric word recognition. *Brain & Cognition*, 59, 314-321.

Basar, E. (2006) The theory of the whole-brain-work. *International Journal of Psychophysiology*, 60, 133-138.

Basar, E., Hari, R., Lopes da Silva, F.H., Schurmann, M. (1997) Brain alpha activity: new aspects and functional correlates. *International Journal of Psychophysiology*. 26, 1 -481.

Bazanov, O.M., Vernon, D. (2013) Interpreting EEG alpha activity. *Neuroscience and Biobehavioural Reviews*, this issue.

Bearden, T.S., Cassisi, J.E., Pineda, M. (2003) Neurofeedback training for a patient with thalamic and cortical infarctions. *Applied Psychophysiology and Biofeedback*, 28, 241-253.

Becerra J, Fernandez T, Roca-Stappung M, Diaz-Comas L, Galan L, Bosch J, et al. (2012) Neurofeedback in healthy elderly human subjects with electroencephalographic risk for cognitive disorder. *Journal of Alzheimer's Disease*, 28, 357-67.

Beatty, J., Greenberg, A., Deibler, W. P., O'Hanlon, J. F. (1974). Operant control of occipital theta rhythm affects performance in a radar monitoring task. *Science*, 183, 871-873.

1 Berman, B.D., Horovitz, S.G., Venkataraman, G., Hallett, H. (2012) Self-modulation of
2 primary motor cortex activity with motor and motor imagery tasks using real-time fMRI-
3 based neurofeedback. *NeuroImage*, 59, 917 – 925.<http://dx.doi.org/10.1016/j>.
4
5
6

7
8 Berner I, Schabus M, Wienerroither T, Klimesch, W. (2006) The significance of sigma
9 neurofeedback training on sleep spindles and aspects of declarative memory. *Applied*
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

114
115
116
117
118
119
120
121
122
123
124
125
126
127
128
129
130
131
132
133
134
135
136
137
138
139
140
141
142
143
144
145
146
147
148
149
150
151
152
153
154
155
156
157
158
159
160
161
162
163
164
165
166
167
168
169
170
171
172
173
174
175
176
177
178
179
180
181
182
183
184
185
186
187
188
189
190
191
192
193
194
195
196
197
198
199
200
201
202
203
204
205
206
207
208
209
210
211
212
213
214
215
216
217
218
219
220
221
222
223
224
225
226
227
228
229
230
231
232
233
234
235
236
237
238
239
240
241
242
243
244
245
246
247
248
249
250
251
252
253
254
255
256
257
258
259
260
261
262
263
264
265
266
267
268
269
270
271
272
273
274
275
276
277
278
279
280
281
282
283
284
285
286
287
288
289
290
291
292
293
294
295
296
297
298
299
300
301
302
303
304
305
306
307
308
309
310
311
312
313
314
315
316
317
318
319
320
321
322
323
324
325
326
327
328
329
330
331
332
333
334
335
336
337
338
339
340
341
342
343
344
345
346
347
348
349
350
351
352
353
354
355
356
357
358
359
360
361
362
363
364
365
366
367
368
369
370
371
372
373
374
375
376
377
378
379
380
381
382
383
384
385
386
387
388
389
390
391
392
393
394
395
396
397
398
399
400
401
402
403
404
405
406
407
408
409
410
411
412
413
414
415
416
417
418
419
420
421
422
423
424
425
426
427
428
429
430
431
432
433
434
435
436
437
438
439
440
441
442
443
444
445
446
447
448
449
450
451
452
453
454
455
456
457
458
459
460
461
462
463
464
465
466
467
468
469
470
471
472
473
474
475
476
477
478
479
480
481
482
483
484
485
486
487
488
489
490
491
492
493
494
495
496
497
498
499
500
501
502
503
504
505
506
507
508
509
510
511
512
513
514
515
516
517
518
519
520
521
522
523
524
525
526
527
528
529
530
531
532
533
534
535
536
537
538
539
540
541
542
543
544
545
546
547
548
549
550
551
552
553
554
555
556
557
558
559
560
561
562
563
564
565
566
567
568
569
570
571
572
573
574
575
576
577
578
579
580
581
582
583
584
585
586
587
588
589
590
591
592
593
594
595
596
597
598
599
600
601
602
603
604
605
606
607
608
609
610
611
612
613
614
615
616
617
618
619
620
621
622
623
624
625
626
627
628
629
630
631
632
633
634
635
636
637
638
639
640
641
642
643
644
645
646
647
648
649
650
651
652
653
654
655
656
657
658
659
660
661
662
663
664
665
666
667
668
669
670
671
672
673
674
675
676
677
678
679
680
681
682
683
684
685
686
687
688
689
690
691
692
693
694
695
696
697
698
699
700
701
702
703
704
705
706
707
708
709
710
711
712
713
714
715
716
717
718
719
720
721
722
723
724
725
726
727
728
729
730
731
732
733
734
735
736
737
738
739
740
741
742
743
744
745
746
747
748
749
750
751
752
753
754
755
756
757
758
759
760
761
762
763
764
765
766
767
768
769
770
771
772
773
774
775
776
777
778
779
780
781
782
783
784
785
786
787
788
789
790
791
792
793
794
795
796
797
798
799
800
801
802
803
804
805
806
807
808
809
810
811
812
813
814
815
816
817
818
819
820
821
822
823
824
825
826
827
828
829
830
831
832
833
834
835
836
837
838
839
840
841
842
843
844
845
846
847
848
849
850
851
852
853
854
855
856
857
858
859
860
861
862
863
864
865
866
867
868
869
870
871
872
873
874
875
876
877
878
879
880
881
882
883
884
885
886
887
888
889
890
891
892
893
894
895
896
897
898
899
900
901
902
903
904
905
906
907
908
909
910
911
912
913
914
915
916
917
918
919
920
921
922
923
924
925
926
927
928
929
930
931
932
933
934
935
936
937
938
939
940
941
942
943
944
945
946
947
948
949
950
951
952
953
954
955
956
957
958
959
960
961
962
963
964
965
966
967
968
969
970
971
972
973
974
975
976
977
978
979
980
981
982
983
984
985
986
987
988
989
990
991
992
993
994
995
996
997
998
999
1000
1001
1002
1003
1004
1005
1006
1007
1008
1009
1010
1011
1012
1013
1014
1015
1016
1017
1018
1019
1020
1021
1022
1023
1024
1025
1026
1027
1028
1029
1030
1031
1032
1033
1034
1035
1036
1037
1038
1039
1040
1041
1042
1043
1044
1045
1046
1047
1048
1049
1050
1051
1052
1053
1054
1055
1056
1057
1058
1059
1060
1061
1062
1063
1064
1065
1066
1067
1068
1069
1070
1071
1072
1073
1074
1075
1076
1077
1078
1079
1080
1081
1082
1083
1084
1085
1086
1087
1088
1089
1090
1091
1092
1093
1094
1095
1096
1097
1098
1099
1100
1101
1102
1103
1104
1105
1106
1107
1108
1109
1110
1111
1112
1113
1114
1115
1116
1117
1118
1119
1120
1121
1122
1123
1124
1125
1126
1127
1128
1129
1130
1131
1132
1133
1134
1135
1136
1137
1138
1139
1140
1141
1142
1143
1144
1145
1146
1147
1148
1149
1150
1151
1152
1153
1154
1155
1156
1157
1158
1159
1160
1161
1162
1163
1164
1165
1166
1167
1168
1169
1170
1171
1172
1173
1174
1175
1176
1177
1178
1179
1180
1181
1182
1183
1184
1185
1186
1187
1188
1189
1190
1191
1192
1193
1194
1195
1196
1197
1198
1199
1200
1201
1202
1203
1204
1205
1206
1207
1208
1209
1210
1211
1212
1213
1214
1215
1216
1217
1218
1219
1220
1221
1222
1223
1224
1225
1226
1227
1228
1229
1230
1231
1232
1233
1234
1235
1236
1237
1238
1239
1240
1241
1242
1243
1244
1245
1246
1247
1248
1249
1250
1251
1252
1253
1254
1255
1256
1257
1258
1259
1260
1261
1262
1263
1264
1265
1266
1267
1268
1269
1270
1271
1272
1273
1274
1275
1276
1277
1278
1279
1280
1281
1282
1283
1284
1285
1286
1287
1288
1289
1290
1291
1292
1293
1294
1295
1296
1297
1298
1299
1300
1301
1302
1303
1304
1305
1306
1307
1308
1309
1310
1311
1312
1313
1314
1315
1316
1317
1318
1319
1320
1321
1322
1323
1324
1325
1326
1327
1328
1329
1330
1331
1332
1333
1334
1335
1336
1337
1338
1339
1340
1341
1342
1343
1344
1345
1346
1347
1348
1349
1350
1351
1352
1353
1354
1355
1356
1357
1358
1359
1360
1361
1362
1363
1364
1365
1366
1367
1368
1369
1370
1371
1372
1373
1374
1375
1376
1377
1378
1379
1380
1381
1382
1383
1384
1385
1386
1387
1388
1389
1390
1391
1392
1393
1394
1395
1396
1397
1398
1399
1400
1401
1402
1403
1404
1405
1406
1407
1408
1409
1410
1411
1412
1413
1414
1415
1416
1417
1418
1419
1420
1421
1422
1423
1424
1425
1426
1427
1428
1429
1430
1431
1432
1433
1434
1435
1436
1437
1438
1439
1440
1441
1442
1443
1444
1445
1446
1447
1448
1449
1450
1451
1452
1453
1454
1455
1456
1457
1458
1459
1460
1461
1462
1463
1464
1465
1466
1467
1468
1469
1470
1471
1472
1473
1474
1475
1476
1477
1478
1479
1480
1481
1482
1483
1484
1485
1486
1487
1488
1489
1490
1491
1492
1493
1494
1495
1496
1497
1498
1499
1500
1501
1502
1503
1504
1505
1506
1507
1508
1509
1510
1511
1512
1513
1514
1515
1516
1517
1518
1519
1520
1521
1522
1523
1524
1525
1526
1527
1528
1529
1530
1531
1532
1533
1534
1535
1536
1537
1538
1539
1540
1541
1542
1543
1544
1545
1546
1547
1548
1549
1550
1551
1552
1553
1554
1555
1556
1557
1558
1559
1560
1561
1562
1563
1564
1565
1566
1567
1568
1569
1570
1571
1572
1573
1574
1575
1576
1577
1578
1579
1580
1581
1582
1583
1584
1585
1586
1587
1588
1589
1590
1591
1592
1593
1594
1595
1596
1597
1598
1599
1600
1601
1602
1603
1604
1605
1606
1607
1608
1609
1610
1611
1612
1613
1614
1615
1616
1617
1618
1619
1620
1621
1622
1623
1624
1625
1626
1627
1628
1629
1630
1631
1632
1633
1634
1635
1636
1637
1638
1639
1640
1641
1642
1643
1644
1645
1646
1647
1648
1649
1650
1651
1652
1653
1654
1655
1656
1657
1658
1659
1660
1661
1662
1663
1664
1665
1666
1667
1668
1669
1670
1671
1672
1673
1674
1675
1676
1677
1678
1679
1680
1681
1682
1683
1684
1685
1686
1687
1688
1689
1690
1691
1692
1693
1694
1695
1696
1697
1698
1699
1700
1701
1702
1703
1704
1705
1706
1707
1708
1709
1710
1711
1712
1713
1714
1715
1716
1717
1718
1719
1720
1721
1722
1723
1724
1725
1726
1727
1728
1729
1730
1731
1732
1733
1734
1735
1736
1737
1738
1739
1740
1741
1742
1743
1744
1745
1746
1747
1748
1749
1750
1751
1752
1753
1754
1755
1756
1757
1758
1759
1760
1761
1762
1763
1764
1765
1766
1767
1768
1769
1770
1771
1772
1773
1774
1775
1776
1777
1778
1779
1780
1781
1782
1783
1784
1785
1786
1787
1788
1789
1790
1791
1792
1793
1794
1795
1796
1797
1798
1799
1800
1801
1802
1803
1804
1805
1806
1807
1808
1809
1810
1811
1812
1813
1814
1815
1816
1817
1818
1819
1820
1821
1822
1823
1824
1825
1826
1827
1828
1829
1830
1831
1832
1833
1834
1835
1836
1837
1838
1839
1840
1841
1842
1843
1844
1845
1846
1847
1848
1849
1850
1851
1852
1853
1854
1855
1856
1857
1858
1859
1860
1861
1862
1863
1864
1865
1866
1867
1868
1869
1870
1871
1872
1873
1874
1875
1876
1877
1878
1879
1880
1881
1882
1883
1884
1885
1886
1887
1888
1889
1890
1891
1892
1893
1894
1895
1896
1897
1898
1899
1900
1901
1902
1903
1904
1905
1906
1907
1908
1909
1910
1911
1912
1913
1914
1915
1916
1917
1918
1919
1920
1921
1922
1923
1924
1925
1926
1927
1928
1929
1930
1931
1932
1933
1934
1935
1936
1937
1938
1939
1940
1941
1942
1943
1944
1945
1946
1947
1948
1949
1950
1951
1952
1953
1954
1955
1956
1957
1958
1959
1960
1961
1962
1963
1964
1965
1966
1967
1968
1969
1970
1971
1972
1973
1974
1975
1976
1977
1978
1979
1980
1981
1982
1983
1984
1985
1986
1987
1988
1989
1990
1991
1992
1993
1994
1995
1996
1997
1998
1999
2000
2001
2002
2003
2004
2005
2006
2007
2008
2009
2010
2011
2012
2013
2014
2015
2016
2017
2018
2019
2020
2021
2022
2023
2024
2025
2026
2027
2028
2029
2030
2031
2032
2033
2034
2035
2036
2037
2038
2039
2040
2041
2042
2043
2044
2045
2046
2047
2048
2049
2050
2051
2052
2053
2054
2055
2056
2057
2058
2059
2060
2061
2062
2063
2064
2065
2066
2067
2068
2069
2070
2071
2072
2073
2074
2075
2076
2077
2078
2079
2080
2081
2082
2083
2084
2085
2086
2087
2088
2089
2090
2091
2092
2093
2094
2095
2096
2097
2098
2099
2100
2101
2102
2103
2104
2105
2106
2107
2108
2109
2110
2111
2112
2113
2114
2115
2116
2117
2118
2119
2120
2121
2122
2123
2124
2125
2126
2127
2128
2129
2130
2131
2132
2133
2134
2135
2136
2137
2138
2139
2140
2141
2142
2143
2144
2145
2146
2147
2148
2149
2150
2151
2152
2153
2154
2155
2156
2157
2158
2159
2160
2161
2162
2163
2164
2165
2166
2167
2168
2169
2170
2171
2172
2173
2174
2175
2176
2177
2178
2179
2180
2181
2182
2183
2184
2185
2186
2187
2188
2189
2190
2191
2192
2193
2194
2195
2196
2197
2198
2199
2200
2201
2202
2203
2204
2205
2206
2207
2208
2209
2210
2211
2212
2213
2214
2215
2216
2217
2218
2219
2220
2221
2222
2223
2224
2225
2226
2227
2228
2229
2230
2231
2232
2233
2234
2235
2236
2237
2238
2239
2240

1 Cavanagh, J.F., Zambrano-Vazquez L., Allen, J.B. (2011) Theta lingua franca: A common
2 mid-frontal substrate for action monitoring processes. *Psychophysiology*, 49, 220 – 238.
3
4 DOI: 10.1111/j.1469-8986.2011.01293.
5
6
7 Clark, C.R., Veltmeyer, M.D., Hamilton, R.J., Simms, E., Paul, R., Hermens, B., Gordon, E.
8
9 (2004) Spontaneous alpha peak frequency predicts working memory performance across the
10 age span. *International Journal of Psychophysiology*, 53, 1–9.
11
12
13
14 doi:10.1016/j.ijpsycho.2003.12.011
15
16
17 Coben, R., Linden, M., Myers, T.E. (2010). Neurofeedback for autistic spectrum disorder: A
18 review of the literature. *Applied Psychophysiology and Biofeedback*, 35, 83-105.
19
20
21 Cooper, N.R., Croft, R.J., Dominey, S.J., Burgess, A.P., Gruzelier, J., H. (2003) Paradox lost?
22 Exploring the role of alpha oscillations during externally vs. internally directed attention and
23 the implications for idling and inhibition hypotheses. *International Journal of*
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Duffy, E. (1957) The psychological significance of 'arousal' or 'activation'. *Psychological Review*, 64, 265 – 275.

Duffy, F.H., Albert, M.S., McAnulty, G., Garvey, A.J. (1984) Age-related differences in brain electrical activity of healthy subjects. *Annals Neurology*, 16, 430-8.

Doppelmayr, M., Weber, E. (2011). Effects of SMR and theta/beta neurofeedback on reaction time, spatial abilities and creativity. *Journal Neurotherapy*, 15, 115–129.

Duschek, S., Schuepbach D., Doll, A., Werner, N.S., Reys del Paso, G.A. (2011). Self-regulation of cerebral blood flow by means of transcranial doppler sonography biofeedback. *Annals of Behavioral Medicine*, 41, 235-242.

Edge, J., Lancaster, L. (2004). Phenomenological analysis of superior musical performance facilitated by neurofeedback: Enhancing musical performance through neurofeedback: playing the tune of life. *Transpersonal Psychology Review*, 8, 23–35

Egner, T. & Gruzelier, J. H. (2001). Learned self-regulation of EEG frequency components affects attention and event-related brain potentials in humans. *NeuroReport*, 12, 18, 411-415.

Egner, T., Strawson, E., Gruzelier, J.H. (2002) EEG signature and phenomenology of alpha/theta neurofeedback training versus mock feedback. *Applied Psychophysiology and Biofeedback*, 27, 261-270.

Egner, T. Gruzelier JH (2003) Ecological validity of neurofeedback: Modulation of slow wave EEG enhances musical performance. *NeuroReport*, 14, 1225-1228.

Egner, T. Gruzelier, J.H. (2004a) EEG biofeedback of low beta band components: Frequency-specific effects on variables of attention and event-related brain potentials. *Clinical Neurophysiology*, 115, 131-139.

Egner, T. Gruzelier, J.H. (2004b) The temporal dynamics of electroencephalographic responses to alpha/theta neurofeedback training in healthy subjects. *Journal of Neurotherapy*, 8, 43-57.

1 Egner, T., Zech, T.F., Gruzelier, J.H. (2004) The effects of neurofeedback training on the
2 spectral topography of the healthy electroencephalogram, *Clinical Neurophysiology*, 115,
3 2452-2460.
4
5
6
7 Elbert. T., Rockstroh, B., Lutzenberger, W., Birbaumer, N. (1984) *Self-Regulation of the*
8
9 *Brain and Behavior*. Berlin: Springer.
10
11 Enriques-Geppert, S., Huster, R.J., Herrmann, C.S. (2013) Boosting brain functions: Improving
12 executive functions with behavioral trainings, neurostimulation, and neurofeedback.
13
14
15
16
17 *International Journal of Psychophysiology*, 88, 1-16.
18
19 Enriques-Geppert, S., Huster, R.J., Scharfenort, R., Mokom, Z.N., Zimmermann, J., Herrmann,
20
21 C.S. (2013) Modulation of frontal-midline theta by neurofeedback. *Biological Psychology*,
22
23
24 SAN special issue, in press.
25
26
27 Escolano, C., Aguilar, M., Minguez, J. (2011) EEG-based upper alpha neurofeedback
28
29 training improves working memory performance. 33rd Annual International Conference of
30
31 the IEEE EMBS, Boston, August 30 - September 3.
32
33
34 Fan, J., McCandliss, B.D., Sommer, T., Raz, A., Posner, M.I. (2002). Testing the efficiency and
35
36 independence of attentional networks. *Journal of Cognitive Neuroscience*, 14, 340-347.
37
38
39 Faridnia, M., Shojaei, M., Rahimi, A. (2012) The effect of neurofeedback on the anxiety of
40
41 elite female swimmers. *Annals of Biological Research*, 3, 1020 – 1028.
42
43
44 Feuerstein, M., Labbé, E.E., Kuczmierczyk, A.R. (1988) *Health Psychology: A*
45
46 *Psychobiological Perspective*.
47
48
49 Fitts, P.M., Posner, M.I. (1967) *Human Performance*. Belmont, Brooks/Cole, 1- 162.
50
51
52 Fogel, S. M., Nader, R., Cote, K. A., Smith, C. T. (2007) Sleep spindles and learning
53
54 potential. *Behavioral Neuroscience*, 121, 1-10. doi: 10.1037/0735-7044.121.1.1
55
56
57 Gevensleben, H., Björn, B.H., Vogel, C., Schlamp, D., Kratz, O., Studer, P., Rothenberger, A.,
58
59 Moll, G.H., Heinrich, H. (2009) Is neurofeedback an efficacious treatment for ADHD? A
60
61
62
63
64
65

randomised controlled clinical trial. *Journal of Child Psychology and Psychiatry*, 50, 780–789.

Greenberg, L.M. Kindschi, C.L. (1999) *Test of variables of attention: Clinical guide*.

Universal Attention Disorders, Inc. St Paul, MN.

Gruber, T., Tsivilis, D., Giabbiconi, C.M., Müller, M.M. (2008) Induced

electroencephalogram oscillations during source memory: familiarity is reflected in the

gamma band, recollection in the theta band. *Journal of Cognitive Neuroscience*, 20, 1043–

1053.

Grunwald, M., Weiss, T., Krause, W., Beyer, L., Rost, R., Gutberlet, I., Gertz, H.J. (2001)

Theta power in the EEG of humans during ongoing processing in a haptic object recognition task. *Cognitive Brain Research*, 11, 33-7.

Gruzelier, J.H. (2012) Enhancing imaginative expression in the performing arts with EEG-

neurofeedback. In D. Miell, R. MacDonald, D. Hargreaves, eds. *Musical Imaginations:*

Multidisciplinary perspectives on creativity, performance and perception. Oxford University Press, Oxford, 332-350.

Gruzelier, J. H. (2009) A theory of alpha/theta neurofeedback, creative performance

enhancement, long distance functional connectivity and psychological integration. *Cognitive Processing*, 10, 101-110.

Gruzelier, J.H., (2013a) EEG-neurofeedback for optimising performance II: A review of creativity. *Neuroscience and Biobehavioural Reviews*. SAN special issue, Applied Neuroscience, submitted.

Gruzelier, J.H., (2013b) EEG-neurofeedback for optimising performance III: A review of methodological and theoretical considerations. *Neuroscience and Biobehavioural Reviews*. SAN special issue, Applied Neuroscience, in preparation.

Gruzelier, J.H., (2013c). Differential effects on mood of 12–15 (SMR) and 15–18 (beta1) Hz

neurofeedback. International Journal of Psychophysiology, <http://dx.doi.org/10.1016/j.ijpsycho.2012.11.007>.

Gruzelier, J.H., Egner, T., Vernon, D. (2006) Validating the efficacy of neurofeedback for optimising performance. In C. Neuper, W. Klimesch, Event-related dynamics of brain oscillations. Progress in Brain Research, 159, 421-431.

Gruzelier J.H., Foks M., Steffert, T., Chen, M.J., Ros, T., (2013). Beneficial outcome from EEG-neurofeedback on creative music performance, attention and well-being in school children. Biological Psychology. Apr 25. doi:pii: S0301-0511(13)00099-9. 10.1016/j.

Haarmann, H.J., Davelaar, E.J., Usher, M. (2003) Individual differences in semantic short-term memory capacity and reading comprehension. Journal of Memory and Language, 48, 320 – 345.

Haarmann, H.J. and Cameron, K.A. (2005) Active maintenance of sentence meaning in working memory: Evidence from EEG coherences. 57, 115 – 128.

Hartmann, T., Lorenz, I., Muller, N., Langguth, B., Weisz, N. (2013) The Effects of neurofeedback on oscillatory processes related to tinnitus. Brain Topography.

Hamilton, P. J., Glover, G.H., Hsu, J-J., Johnson, R.F., Gotlib, I.H. (2010) Modulation of subgenual anterior cingulate cortex activity with real-time neurofeedback. Human Brain Mapping, 32, 22–31. DOI: 10.1002/hbm.20997

Hammond, D.C. (2005) Neurofeedback treatment of depression and anxiety. Journal of Adult Development. 12, 131-137.

Hanslmayer, S., Sauseng, P. Doppelmayr, M., Schabus, M., Klimesch, W. (2006) Increasing individual upper alpha by neurofeedback improves cognitive performance in human subjects. Journal of Applied Psychophysiology Biofeedback, 30, 1-10.

Hardt, J.V. and Kamiya, J. (1978). Anxiety change through electroencephalographic alpha

feedback seen only in high anxiety subjects. *Science*, 201, 79-81.

Hartmann T, Lorenz I, Müller N, Langguth B, Weisz N. (2013) The effects of neurofeedback on oscillatory processes related to tinnitus. *Brain Topography* [pub ahead of print] DOI 10.1007/s10548-013-0295-9.

Hauri, P.J. (1981) Treating psychophysiologic insomnia with biofeedback. *Arch Gen Psychiatry*, 38, 752-58.

Hauri, P.J., Percy, L., Hellekson, C., Hartmann, E., Russ, D. (1982) The treatment of psychophysiologic insomnia with biofeedback: a replication study. *Biofeedback and Self-Regulation*, 7, 223-35.

Hoedlmoser, K., Pecherstorfer, T., Gruber, G., Anderer, P., Doppelmayr, M., Klimesch, W., Scgabus, M. (2008) Instrumental conditioning of human sensorimotor rhythm (12-15Hz) and its impact on sleep as well as declarative learning. *Sleep*, 31, 1401-1408.

Hommel, B. (1998). Event files: Evidence for automatic integration of stimulus–response episodes. *Visual Cognition*, 5, 183–216.

Hughes, S.W., Lorincz, M., Cope, D.W., Blethyn, K.L., Kekesi, K.A., Parri, H.R., Juhasz, G., Crunelli, V. (2004) Synchronized oscillations at alpha and theta frequencies in the lateral geniculate nucleus. *Neuron*, 42, 253-268.

Inouye, T., Shinosaki, K., Iyama, A., Matsumoto, Y., Toi, S., Ishihara, T. (1994) Potential flow of frontal midline theta activity during a mental task in the human electroencephalogram. *Neuroscience Letters*, 169, 145 – 148.

Jausovec, N., Jausovec, K. (2000). Differences in resting EEG related to ability. *Brain Topography*, 12, 229–240.

Jaušovec, N., Jaušovec, K., Gerlič, I. (2001) Differences in event-related and induced EEG patterns in the theta and alpha frequency bands related to human emotional intelligence. *Neuroscience Letters*, 311, 93–96.

Jensen, O., Tesche, C.D., (2002) Frontal theta activity in humans increases with memory load in a working memory task. *European Journal of Neuroscience*, 15, 1395 – 1399.

DOI: 10.1046/j.1460-9568.2002.01975.

Kamiya, J. (1968) Conscious control of brain waves. *Psychology Today*, 1, 57–60.

Kamiya, J (1969). Operant control of the EEG alpha rhythm and some of its reported effects on consciousness. In CT Tart (Ed), *Altered states of consciousness*, pp 519-529. Wiley, New York.

Keizer, A.W., Verschoor, M., Verment, R.S., Hommel, B. (2010a) The effect of gamma enhancing neurofeedback on the control of feature bindings and intelligence measures.

International Journal of Psychophysiology, 75, 25-32.

Keizer, A.W., Verment, R.S., Hommel, B. (2010b) Enhancing cognitive control through neurofeedback: A role of gamma-band activity in managing episodic retrieval. *NeuroImage*, 49, 3404-3413.

Klimesch, W. (1999) EEG alpha and theta oscillations reflect cognitive and memory performance: A review and analysis. *Brain Research Reviews*, 29, 169–195.

Klimesch, W., Schimke, H., Pfurtscheller, G. (1993) Alpha frequency, cognitive load and memory performance. *Brain Topography*. 5, 241-251.

Klimesch, W., Doppelmayr, M., Schimke, H., Pachinger, T. (1996) Alpha frequency, reaction time and the speed of processing information. *Journal of Clinical Neurophysiology*, 13, 511-518.

Klimesch, W., Sauseng, P., Hanslmayr, S. (2007). EEG alpha oscillations: The inhibition–timing hypothesis. *Brain Research Review*. 53, 63–88.

Kober, S.E., Wood, G., Våljamäe, A., Stangl, M., Wippel, T., Kurzmann, J., Neuper, C. (2013). Near-Infrared Spectroscopy based neurofeedback training increases specific motor imagery related cortical activation compared to sham feedback. *Biological Psychology*,

SAN special issue Neurofeedback, in press.

Kubota, Y., Sato, W., Toichi, M., Murai, T., Okada, T. (2001) Frontal midline theta rhythm is correlated with cardiac autonomic activities during the performance of an attention demanding meditation procedure. *Cognitive Brain Research*, 11, 281 – 287.

Lagopoulos, J., Xu, J., Rasmussen, I., Vik, A., Mahli, g.S., Eliassen, C.F., Arntsen, I.E., Saether, J.G., et al (2009) Increased theta and alpha EEG activity during nondirective meditation *Journal of Alternative and Complimentary Medicine*, 15, 1187 – 1192.

Landers, D. M., Petruzzello, S. J., Salazar, W., Crews, D. J., Kubitz, K. A., Gannon, T. L., et al. (1991). The influence of electrocrtical biofeedback on performance in pre-elite archers. *Medicine and Science in Sports and Exercise*, 23, 123–129.

Lecomte, G., Juhel, J. (2011) The effects of neurofeedback training on memory performance in elderly subjects. *Psychology*, 2, 846-852.

Leins, U., Goth, G., Hinterberger, T., Klinger, C., Rumpf, N., Strehl, U. (2007) Neurofeedback for children with ADHD: a comparison of SCP and Theta/Beta protocols. *Applied Psychophysiology and Biofeedback*, 32, 73-88.

Levesque, J. Beauregard, M., Mensour, B. (2006) Effect of neurofeedback training on the neural substrates of selective attention in children with attention-deficit/hyperactivity disorder : A functional magnetic resonance imaging study. *Neuroscience Letters*, 394, 216-221.

Linden, M., Habib, T., Radojevic, V. (1996) A controlled study of the effects of EEG biofeedback on cognition and behavior of children with attention deficit disorder and learning disabilities. *Biofeedback and Self-regulation*, 21, 35 – 49.

Lindsley, D. B. (1952) Psychological phenomena and the electroencephalogram. *Electroencephalography and clinical Neurophysiology*, 4, 443-456.

Lofthouse, N., Arnold, L.E., Hersch1, S., Hurt, E., DeBeus, R. (2012) A review of

neurofeedback treatment for pediatric ADHD. *Journal of Attention Disorders* 16, 351-372.

Lopes da Silva, F.H., Pijn, J., Veris, D., Nijssen, P.C.G. (1997) Alpha rhythms: noise, dynamics and models. *International Journal of psychophysiology*, 26, 237 – 250.

Lubar, J.O., Lubar, J.F. (1984) Electroencephalographic biofeedback of SMR and beta for treatment of Attention Deficit Disorders in a clinical setting. *Biofeedback and Self-Regulation*, 9, 1-23.

Lubar, J.F., Shouse, M.N. (1976) EEG and behavioural changes in a hyperkinetic child concurrent with training of the sensorimotor rhythm (SMR): A preliminary report. *Biofeedback and Self-Regulation*, 3, 293-306.

Lubar, J.F., Swartwood, M.O., Swartwood, J.N., O'Donnell, P.H. (1995a). Evaluation of the effectiveness of EEG neurofeedback training for ADHD in a clinical setting as measured by changes in T.O.V.A. scores, behavioural ratings, and WISC-R performance. *Biofeedback and Self-Regulation*, 20, 83–99.

Lubar, J.F., Swartwood, M.O., Swartwood, J.N., Timmerman, D.L. (1995b). Quantitative EEG and auditory event related potentials in the evaluation of attention deficit/hyperactivity disorder: effects of methylphenidate and implications for neurofeedback training. *Journal of Psychoeducational Assessment*, 4, 143–160.

Mak, J.N., Wolpaw, J.R. (2009) Clinical applications of brain-computer interfaces: current state and future prospects. *Biomedical Engineering, IEEE Reviews*, 2, 187 – 199.

Martens, R. (1977) *Sport Competition Anxiety Test*. Champaign, Human Kinetics Publishers.

McNair, D.M., Lorr, D., Droppleman, M.F. (1992) *Profile of Mood States Manual*, Educational and Industrial Testing Service, San Diego.

Mihara, M., Miyai, I., Hattori, N., Hatakenaka, M., Yagura, H., et al. (2012) Neurofeedback using real-time Near-Infrared Spectroscopy enhances motor imagery related cortical activation. *PLoS ONE* 7: e32234. doi:10.1371/journal.pone.0032234.

1 Missonnier, P., Deiber, M-P. (2006) Frontal theta event-related synchronization: comparison
2 of directed attention and working memory load effects. *Journal of Neural Transmission*, 113,
3 1477-1486.
4
5
6
7 Mitchell, D.J., McNaughton, N., Flanagan, D., Kirk, I.J. (2008) Frontal-midline theta from
8 the perspective of hippocampal "theta". *Progress in Neurobiology*, 86, 156-85.
9
10
11
12 Monastra, V. J., Lynn, S., Linden, M., Lubar, J. F., Gruzelier, J., & LaVaque, T. J. (2005).
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Nan, W., Rodrigues, J.P., Ma, J., Qu, X., Wan, F., Mak, P-I., Mak, P.U., Vai, M.I., Rosa, A.
(2012) Individual alpha neurofeedback training effect on short term memory. *International
Journal of Psychophysiology*, 86, 83–87.

Niedermeyer ,E. (2005) The normal EEG of the waking adult. In: Niedermeyer E, Lopes da
Silva F, eds. *Electroencephalography: basic principles, clinical applications, and related
fields*. Baltimore: Williams & Williams, 149-73.

Nissen, N.J., Bullemer, P. (1987) Attentional requirements of learning: Evidence from
performance measures. *Cognitive Psychology*, 19, 1-32.

Onton, J., Delorme, A., Makeig, S. (2005) Frontal midline EEG dynamics during working
memory. *NeuroImage*. 27, 341 – 356

Owen, A.M., Hampshire, A., Grahn, J.A., Stenton, R., Dajani, S. Burns, A.S., Howard, R.J.,
Ballard. C.G. (2010) Putting brain training to the test. *Nature*, 465, 775–778.

Peniston E. G., Kulkosky P. J. (1989). Alpha-theta brainwave training and beta endorphin
levels in alcoholics. *Alcoholism: Clinical and Experimental Research*, 13, 271-279.

Peniston, E. G., & Kulkosky, P. J. (1991). Alpha-theta brainwave neurofeedback for
Vietnam veterans with combat-related post-traumatic stress disorder. *Medical Psychotherapy*,
4, 47-60.

Petsche, H., Kaplan, S., von Stein, A., Filz, O. (1997). The possible meaning of the upper and lower alpha frequency ranges for cognitive and creative tasks. *International Journal of Psychophysiology*, 26, 77-97.

Posner, M.I., Raichle, M.E. (1996). *Images of Mind*, Review. Washington, DC: Scientific American Books.

Posner, M.I., Peterson, S.E. (1990) The attention systems of the human brain. *Annual Review of Neuroscience*. 19, 25–42.

Ray, W.J., Cole, H.W. (1985) EEG alpha activity reflects attentional demands, and beta activity reflects emotional and cognitive processes. *Science*, 228, 750 – 752. DOI: 10.1126/science.3992243

Raymond, J. Varney, C. Gruzelier J.H. (2005) The effects of alpha/theta neurofeedback on personality and mood. *Cognitive Brain Research*, 23, 287-292.

Rabipour, S. Raz, A. (2012) Training the brain: fact or fad in cognitive and behavioural remediation. *Brain and Cognition*, 79, 159 -179.

Rauchs, G., Desgranges, B., Foret, J., Eustache, F. (2005) The relationships between memory systems and sleep stages. *Journal of Sleep Research*, 14, 123-140.

Raven, J.C. (1938). *Progressive Matrices: A Perceptual Test of Intelligence*. H.K. Lewis, London.

Reiner, M., Rozengurt, R., Barnea, A. (2013) Theta neurofeedback training and enhancement of motor performance. *Biological Psychology*, Neurofeedback special issue under revision.

Rockstroh, B. (1989) *Slow Cortical Potentials and Behaviour*. Urban & Schwarzenberg

Ros, T., Moseley, M.J., Bloom, P.A., Benjamin, L. Parkinson, L.A., Gruzelier, J.H. (2009) Optimizing microsurgical skills with EEG-neurofeedback. *BMC Neuroscience*, 10, 87
doi:10.1186/1471-2202-10-87

Ros, T., Munneke, M.A.M., Ruge, D., Gruzelier, J.H., Rothwell, J.C. (2010) Endogenous

control of waking alpha rhythms induces neuroplasticity. *European Journal of Neuroscience*, 31, 770-778.

Ros, T., Munneke, M.A.M., Parkinson, L. A., Gruzelier, J.H. (2013) Facilitation of perceptuo-motor learning with one session of neurofeedback. *Biological Psychology*, neurofeedback special issue in press.

Rossiter, TR, LaVaque, TJ (1995). A comparison of EEG biofeedback and psychostimulants in treating Attention Deficit Hyperactivity Disorders. *Journal of Neurotherapy*, 1, 48-59.

Rota, G., Sitaram, R., Veit R., Erb, M., Weiskopf, N., Dogil, D., Birbaumer, N., 2009. Self-regulation of regional cortical activity using real-time fMRI: The right inferior frontal gyrus and linguistic processing. *Human Brain Mapping* 30, 1605 – 14.

Ruiz, S., Sitaram, R., Birbaumer, N., 2013 Real-time fMRI Brain-Computer Interfaces: Operant training of single brain regions to networks. *Biol Psychol*, special issue Neurofeedback, in press.

Sarnthein. J., Petsche, H., Rappelsberger, P., Shaw, G.L., von Stein, A. (1998) Synchronization between prefrontal and posterior association cortex during human working memory. *Science*, 95, 7092 – 7096.

Saxby, E., & Peniston, E. G. (1995). Alpha-theta brainwave neurofeedback training: An effective treatment for male and female alcoholics with depressive symptoms. *Journal of Clinical Psychology*, 51, 685-693.

Schabus M, Gruber G, Parapatics S, et al. (2004) Sleep spindles and their significance for declarative memory consolidation. *Sleep*, 27,1479-85.

Schabus, M., Heib, D.P.J., Lechinger, J., Griessenberger, H., Klimesch, W., Pawlizki, A., Kunz, A.B., Sterman, B.M., Hoedlmoser, K. (2013) Enhancing sleep quality and memory in insomnia using instrumental sensorimotor rhythm conditioning. *Biological psychology*, neurofeedback special issue in press.

Silvoni, S., Ramos-Murguialday, A., Cavinato, M. Volpato, C., Cisotto, G., Turolla, A.,
 Piccione, F., Birbaumer, N. (2011) Brain-computer interface in stroke: A review of
 progress. *Clinical EEG and Neuroscience*. 42, 245-252.

Song, S. (2009) Consciousness and the consolidation of motor learning. *Behavioural
 Brain Research*, 196, 180-186.

Staufenbiel, S.W., Brouwer, A.M., Keizer, A.W., Van Wouwe, N.C. (2013) Effect of beta
 and gamma neurofeedback on memory and intelligence in the elderly. *Biological Psychology*,
 Neurofeedback special issue in press.

Stam, C.J., Van Walsum, A.M.V., Pijnenburg, Y.A.L., Berendse, H.W., de Munck, J.C.,
 Scheltens, P., et al. (2002). Generalized synchronization of MEG recordings in
 Alzheimer's Disease: Evidence for involvement of the gamma band. *Journal of
 Clinical Neurophysiology*, 19(6), 562-574.

Sterman, M.B. (1996). Physiological origins and functional correlates of EEG rhythmic
 activities: Implications for self-regulation. *Biofeedback and Self-Regulation*, 21, 3-33.

Sterman, M. B. (2000). Basic concepts and clinical findings in the treatment of seizure
 disorders with EEG operant conditioning. *Clinical Electroencephalography*, 31, 35-45.

Sterman, M.B., Wyrwicka, W. (1967). EEG correlates of sleep: Evidence for separate
 forebrain substrates. *Brain Research*, 6, 143-163.

Sterman, M.B. and Friar, L. (1972). Suppression of seizures in an epileptic following
 sensorimotor EEG feedback training. *Electroencephalography and Clinical Neurophysiology*,
 33, 89-95.

Sterman, M.B., Wyrwicka, W., Roth, S.R. (1969). Electrophysiological correlates and neural
 substrates of alimentary behaviour in the cat. *Annals of the New York Academy of Science*,
 157, 723-739.

Sterman, M.B., Howe, R.C., MacDonald, L.R. (1970) Facilitation of spindle-burst sleep by

conditioning of electroencephalographic activity while awake. *Science*. 167,1146-48.

Sterman, M.B., MacDonald, L.R., Stone, R.K. (1974). Biofeedback training of the sensorimotor EEG rhythm in man: Effects on epilepsy. *Epilepsia*, 15, 395-417.

Sterman, M.B., MacDonald, L.R. (1978). Effects of central cortical EEG feedback training on incidence of poorly controlled seizures. *Epilepsia*, 19, 207-222.

Sternberg, S. (1966) High-speed scanning in human memory. *Science*. 153, 652-4.

Tan, G., Thornby, J., Hammond, D.C., Strehl, U., Canady, B., Arnemann, K., Kaiser, D.A. (2009) Meta-analysis of EEG biofeedback in treating epilepsy. *Clinical EEG and Neuroscience*. 40, 173-179.

Thayer, R., E. (1967) Measurement of activation through self-report. *Psychological Reports*. 20, 663–678.

Thompson, T., Steffert, T., Ros, T., Leach, J., Gruzelier, J. (2008) EEG applications for sport and performance. *Methods*, 45, 279-288.

Unterrainer, H, Chen, M. J-L, Gruzelier, J.H., (2013) EEG-neurofeedback and psychodynamic psychotherapy in adolescent anhedonia with substance misuse: A single case study. *International Journal of Psychophysiology*, doi:pii: S0167-8760(13)00061-5. 10.1016/j.ijpsycho.2013.03.011.

Van Boxtel, G.J.M., Denissen, A., Jager, M., Vernon, D., Dekker, M.K.J., Mihaljovic, V. & Sitskoorn, M.M. (2012). A novel self-guided approach to alpha activity training. *International journal of psychophysiology*, 83, 282-294.

Varela, F., Lachaux, J.-P., Rodriguez, E., Martinerie, J. (2001) The brainweb: phase synchronization and large-scale integration. *Nature Review Neuroscience*, 2, 229–239.

Vernon, D., Egner, T., Cooper, N., Compton, T., Neilands, C., Sheri, A., Gruzelier, J. (2003) The effect of training distinct neurofeedback protocols on aspects of cognitive performance. *International Journal of Psychophysiology*, 47, 75-86.

von Stein A., Sarntheim., J. (2000) Different frequencies for different scales of cortical
 integration: from local gamma to long range alpha/theta synchronisation. *International
 Journal of Psychophysiology*, 38, 301–313.

von Stein, A., Chiang, C., Konig, P. (2000) Top-down processing mediated by interareal
 synchronization. *Proceedings National Academy of Sciences USA*, 97, 14748 –14753.

Walker, M. P. (2005). A refined model of sleep and the time course of memory formation." *Behavioural and Brain Sciences*, 28, 51-64.

Wang, J.R., Hsieh, S. (2013) Neurofeedback training improves attention and working
 memory performance. *Clinical Neurophysiology*, [http://dx.doi.org/10.1016/
 j.clinph.2013.05.020](http://dx.doi.org/10.1016/j.clinph.2013.05.020)

Weiskopf, N., Scharnowski, F., Veit, R., Goebel, R., Birbaumer, N.& Mathiak, K. (2004)
 Self-regulation of local brain activity using real-time functional magnetic resonance imaging
 (fMRI). *Journal Physiology Paris*, 98, 357-373.

Werkle-Bergner, M., Shing, Y.L., Müller, V., Li, S.-C., & Lindenberger, U. (2009). EEG
 gamma-band synchronization in visual coding from childhood to old age: Evidence from
 evoked power and inter-trial phase locking. *Clinical Neurophysiology*, 120, 1291-1302.

Wyrwicka, W, Sterman, MB (1968). Instrumental conditioning of sensorimotor cortex EEG
 spindles in the waking cat. *Physiology and Behaviour*, 3, 703-707.

Yoo, S-S., Fairney, T., Chen, N-K., Choo, S-E., Panych, L. P., Park, H-W., Lee, S-Y. &
 Jolesz, F.A. (2004) Brain-computer interface using fMRI: spatial navigation by thoughts
Neuroreport, 15, 1591-1595.

Zoefel , B., Huster , R.J., Hermann , C.S. (2011). Neurofeedback training of the upper alpha
 frequency band EEG improves cognitive performance. *Neuroimage*, 54, 1427 – 31.

1 Zotev, V., Krueger, F., Phillips, R., Alvarez, R.P., Simmons, W.K., et al. (2011) Self-
2 regulation of amygdala activation using real-time fMRI neurofeedback. PLoS ONE 6:
3 e24522. doi:10.1371/journal.pone.0024522
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23

24 Figure Captions

25
26
27
28
29 Figure 1. SMR ratio learning indices positively correlated with the reduction in commission
30 errors in sustained attention ($r = -0.79$, $p < 0.001$) (Egner and Gruzelier, 2001).
31

32
33
34 Figure 2. The increase in amplitude of P300b correlated with learning indices of SMR ratio
35 ($r = 0.49$, $p < 0.06$) and beta1 ratio ($r = 0.55$, $p < 0.05$) protocols (Egner and Gruzelier, 2001).
36

37
38
39 Figure 3. Temporal changes in technical skills in a simulated cataract operation following
40 alpha/theta (A/T) or SMR ratio protocols and in non-intervention controls (C) (Ros et al,
41
42
43 2009).
44

45
46 Figure 4. Improvements in visuospatial rotation in SMR, theta/beta ratio (TBR) and random
47
48
49 bin feedback (TBR) control groups (Dopplemayer and Weber, 2011).
50

51
52 Figure 5. Positive correlations between SMR learning and increase in frequency of fast
53
54
55 spindles (left) ($r = 0.38$, $p < 0.04$) and memory consolidation (right) ($r = 0.40$, $p < 0.027$)
56 (Schabus et al, 2013).
57

58
59 Figure 6. Percentage change in spatial rotation task following one session of upper-alpha
60
61
62
63
64
65

training ($r = 0.41 - 0.55$, $p < 0.05$) (Hanslmayr et al, 2006).

Figure 7. Percentage change in gamma power correlated with percentage change in fluid intelligence ($r = 0.81$, $p < 0.03$) (Keizer et al, 2010a).

Figure 8. Pre and post –training alerting, orienting and conflict scores in elderly (O) and students (Y) receiving frontal theta up-training or sham training (Wang and Hsieh, 2013).

Figure
1.




Figure 2




Figure 3




Figure
4





Figure
5



Figure




Figure 6

Figure




Figure 7




Figure 8